

# American Mathematical Society

Jennifer Wright Sharp  
American Mathematical Society  
201 Charles Street  
Providence, RI 02904  
jws@ams.org

Job ID: pubp-99336

Title: /ams/production/journals/bull/inprod/bull1413/bull1413.dvi

User: jws


Date: 03 Apr 2013 09:47:57

In order to maintain consistency of style in  
this publication, electronic submissions are  
run through our production system.  
Please proofread galleys.

Author: Your references were processed using  
our Enhanced Reference Program. Please see  
<http://www.ams.org/publications/journals/references>  
for information.



**AMS**



BULLETIN (New Series) OF THE  
 AMERICAN MATHEMATICAL SOCIETY  
 S 0273-0979(XX)0000-0

*Introduction to Vassiliev knot invariants*, by S. Chmutov, S. Duzhin, and J. Mostovoy, Cambridge University Press, Cambridge, 2012, xvi+504 pp., ISBN 978-1-107-02083-2, hardcover, US \$70.00.

Merely 30 years ago, if you had asked even the best informed mathematician about the relationship between knots and Lie algebras, she would have laughed, for there isn't and there can't be. Knots are flexible; Lie algebras are rigid. Knots are irregular; Lie algebras are symmetric. The list of knots is a lengthy mess; the collection of Lie algebras is well organized. Knots are useful for sailors, scouts, and hangmen; Lie algebras for navigators, engineers, and high energy physicists. Knots are blue collar; Lie algebras are white. They are as similar as worms and crystals: both well studied, but hardly ever together. Then in the 1980s came Jones,




FIGURE 1. A knot and a Lie algebra, a list of knots and a list of Lie algebras, and an unusual conference of the symmetric and the knotted.

and Witten, and Reshetikhin and Turaev [8, 12, 15] and showed that if you really are the best informed, and you know your quantum field theory and conformal field theory and quantum groups, then you know that the two disjoint fields are in fact intricately related. This “quantum” approach remains the most powerful way to get computable knot invariants out of (certain) Lie algebras (and representations thereof). Yet shortly later, in the late 1980s and early 1990s, an alternative perspective aroused, that of “finite-type” or “Vassiliev–Goussarov” invariants [1, 4, 6, 7, 9, 10, 13, 14], which made the surprising relationship between knots and Lie algebras appear simple and almost inevitable.

2010 *Mathematics Subject Classification*. Primary 57M25.

Picture credits: Rope from “The Project Gutenberg eBook, Knots, Splices and Rope Work, by A. Hyatt Verrill”, <http://www.gutenberg.org/files/13510/13510-h/13510-h.htm>. Plane from NASA, <http://www.grc.nasa.gov/WWW/k-12/airplane/rotations.html>.

TeX at <http://drorbn.net/AcademicPensieve/2013-01/CDMReview/>, copyleft at <http://www.math.toronto.edu/~drorbn/Copyleft/>. This review was written while I was a guest at the Newton Institute, in Cambridge, UK..

The book under review [3] is about that alternative perspective, the one reasonable sounding but not entirely trivial theorem that is crucially needed within it (the *Fundamental Theorem* or the *Kontsevich integral*), and the many threads that begin with that perspective. Let me start with a brief summary of the mathematics, and even before, an even briefer summary.

The briefer summary is that in some combinatorial sense it is possible to “differentiate” knot invariants, and hence it makes sense to talk about “polynomials” on the space of knots — these are functions on the set of knots (namely, these are knot invariants) whose sufficiently high derivatives vanish. Such polynomials can be fairly conjectured to separate knots — elsewhere in mathematics in lucky cases polynomials separate points, and in our case, specific computations are encouraging. Also, such polynomials are determined by their “coefficients”, and each of these, by the one-side-easy Fundamental Theorem, is a linear functional on some finite space of graphs modulo relations. These same graphs turn out to parameterize formulas that make sense in a wide class of Lie algebras, and the said relations match exactly with the relations in the definition of a Lie algebra — anti-symmetry and the Jacobi identity. Hence what is more or less dual to knots (invariants), is also, after passing to the coefficients, dual to certain graphs which are more or less dual to Lie algebras. QED, and on to the less brief summary.<sup>1</sup>

Let  $V$  be an arbitrary invariant of oriented knots in oriented space with values in (say)  $\mathbb{Q}$ . Extend  $V$  to be an invariant of 1-singular knots, knots that have a single singularity that locally looks like a double point  $\times$ , using the formula

$$(1) \quad V(\times) = V(\nearrow) - V(\searrow).$$

Further extend  $V$  to the set  $\mathcal{K}^m$  of  $m$ -singular knots (knots with  $m$  such double points) by repeatedly using (1).

**Definition 1.** We say that  $V$  is of type  $m$  (or “Vassiliev of type  $m$ ”) if its extension  $V|_{\mathcal{K}^{m+1}}$  to  $(m+1)$ -singular knots vanishes identically. We say that  $V$  is of finite type (or “Vassiliev”) if it is of type  $m$  for some  $m$ .

Repeated differences are similar to repeated derivatives and hence it is fair to think of the definition of  $V|_{\mathcal{K}^m}$  as repeated differentiation. With this in mind, the above definition imitates the definition of polynomials of degree  $m$ . Hence finite type invariants can be thought of as “polynomials” on the space of knots.<sup>2</sup> It is known (see, e.g., [3]) that the class of finite type invariants is large and powerful. Yet the first question on finite type invariants remains unanswered:

**Problem 2.** Honest polynomials are dense in the space of functions. Are finite type invariants dense within the space of all knot invariants? Do they separate knots?

The top derivatives of a multi-variable polynomial form a system of constants that determine that polynomial up to polynomials of lower degree. Likewise, the  $m$ th derivative<sup>3</sup>  $V^{(m)} = V|_{\mathcal{K}^m} = V(\times^{\cdot m} \times)$  of a type  $m$  invariant  $V$  is a constant

<sup>1</sup>Partially self-plagiarized from [2].

<sup>2</sup>Keep this apart from invariants of knots whose values are polynomials, such as the Alexander or the Jones polynomial. A posteriori related, these are a priori entirely different.

<sup>3</sup>As common in the knot theory literature, in the formulas that follow a picture such as  $\times^{\cdot m} \times$  indicates “some knot having  $m$  double points and a further (right-handed) crossing”. Furthermore,





FIGURE 2


in the sense that it does not see the difference between overcrossings and undercrossings, and so it is blind to 3D topology. Indeed,  $V(\overline{\text{X}}^m \cdot \overline{\text{X}}^m) - V(\overline{\text{X}}^m \cdot \overline{\text{X}}^m) = V(\overline{\text{X}}^{m+1}) = 0$ . Also, clearly  $V^{(m)}$  determines  $V$  up to invariants of lower type. Hence a primary tool in the study of finite type invariants is the study of the “top derivative”  $V^{(m)}$ , also known as “the weight system of  $V$ ”. Blind to 3D topology,  $V^{(m)}$  only sees the combinatorics of the circle that parameterizes an  $m$ -singular knot. On this circle there are  $m$  pairs of points that are pairwise identified in the image; standardly one indicates those by drawing a circle with  $m$  chords marked (an “ $m$ -chord diagram”); see Figure 2. Let  $\mathcal{D}_m$  denote the space of all formal linear combinations with rational coefficients of  $m$ -chord diagrams. Thus  $V^{(m)}$  is a linear functional on  $\mathcal{D}_m$ .

I leave it for the reader to figure out or read in [3, p. 88] how the following figure easily implies the “4T” relations of the “easy side” of the theorem that follows:



**Theorem 3** (The Fundamental Theorem, details in [3]).


- (Easy side) If  $V$  is a rational valued type  $m$  invariant then  $V^{(m)}$  satisfies the “4T” relations shown below, and hence it descends to a linear functional on  $\mathcal{A}_m := \mathcal{D}_m/4T$ . If in addition  $V^{(m)} \equiv 0$ , then  $V$  is of type  $m - 1$ .



- (Hard side, slightly misstated by avoiding “framings”) For any linear functional  $W$  on  $\mathcal{A}_m$  there is a rational valued type  $m$  invariant  $V$  so that  $V^{(m)} = W$ .

Thus to a large extent the study of finite type invariants is reduced to the finite (though super-exponential in  $m$ ) algebraic study of  $\mathcal{A}_m$ .

when two such pictures appear within the same formula, it is to be understood that the parts of the knots (or diagrams) involved *outside* of the displayed pictures are to be taken as the same.



A Jacobi diagram in a circle





FIGURE 3

Much of the richness of finite type invariants stems from their relationship with Lie algebras. Theorem 4 below suggests this relationship on an abstract level, and Theorem 5 makes that relationship concrete.

**Theorem 4** ([1]). *The space  $\mathcal{A}_m$  is isomorphic to the space  $\mathcal{A}_m^t$  generated by “Jacobi diagrams in a circle” (chord diagrams that are also allowed to have oriented internal trivalent vertices) that have exactly  $2m$  vertices, modulo the AS, STU, and IHX relations; see Figure 3.*

The key to the proof of Theorem 4 is the figure below, which shows that the 4T relation is a consequence of two STU relations. The rest is more or less an exercise in induction.




Thinking of internal trivalent vertices as graphical analogs of the Lie bracket, the AS relation becomes the anti-commutativity of the bracket, STU becomes the equation  $[x, y] = xy - yx$ , and IHX becomes the Jacobi identity. This analogy is made concrete within the following construction, originally due to Penrose [11] and to Cvitanović [5]. Given a finite dimensional metrized Lie algebra  $\mathfrak{g}$  (e.g., any semi-simple Lie algebra) and a finite-dimension representation  $\rho : \mathfrak{g} \rightarrow \text{End}(V)$  of  $\mathfrak{g}$ , choose an orthonormal basis<sup>4</sup>  $\{X_a\}_{a=1}^{\dim \mathfrak{g}}$  of  $\mathfrak{g}$  and some basis  $\{v_\alpha\}_{\alpha=1}^{\dim V}$  of  $V$ , let  $f_{abc}$  and  $r_{\alpha\beta}^\gamma$  be the “structure constants” defined by

$$f_{abc} := \langle [X_a, X_b], X_c \rangle \quad \text{and} \quad \rho(X_a)(v_\beta) = \sum_\gamma r_{\alpha\beta}^\gamma v_\gamma.$$

Now given a Jacobi diagram  $D$  label its circle-arcs with Greek letters  $\alpha, \beta, \dots$ , and its chords with Latin letters  $a, b, \dots$ , and map it to a sum as suggested by the following example:

<sup>4</sup>This requirement can easily be relaxed.



$$\longrightarrow \sum_{a,b,c,\alpha,\beta,\gamma} f_{abc} r_{a\gamma}^{\beta} r_{b\alpha}^{\gamma} r_{c\beta}^{\alpha} \quad \left( \begin{array}{l} \text{internal vertices go to } f\text{'s,} \\ \text{circle-vertices to } r\text{'s} \end{array} \right).$$

**Theorem 5.** *This construction is well defined, and the basic properties of Lie algebras imply that it respects the AS, STU, and IHX relations. Therefore it defines a linear functional  $W_{\mathfrak{g},R} : \mathcal{A}_m \rightarrow \mathbb{Q}$ , for any  $m$ .*

The last assertion along with Theorem 3 show that associated with any  $\mathfrak{g}$ ,  $R$  and  $m$  there is a weight system and hence a knot invariant. Thus knots are indeed linked with Lie algebras.

The above is of course merely a sketch of the beginning of a long story. You can read the details, and some of the rest, in [3].



**What I like about [3].** Detailed, well thought out, and carefully written. Lots of pictures! Many excellent exercises! A complete discussion of “the algebra of chord diagrams”. A nice discussion of the pairing of diagrams with Lie algebras, including examples aplenty. The discussion of the Kontsevich integral (meaning, the proof of the hard side of Theorem 3) is terrific — detailed and complete and full of pictures and examples, adding a great deal to the original sources. The subject of “associators” is huge and worthy of its own book(s); yet in as much as they are related to Vassiliev invariants, the discussion in [3] is excellent. A great many further topics are touched — multiple  $\zeta$ -values, the relationship of the Hopf link with the Duflo isomorphism, intersection graphs and other combinatorial aspects of chord diagrams, Rozansky’s rationality conjecture, the Melvin–Morton conjecture, braids,  $n$ -equivalence, etc.

For all these, I’d certainly recommend [3] to any newcomer to the subject of knot theory, starting with my own students.

However, some proofs other than that of Theorem 3 are repeated as they appear in original articles with only a superficial touch-up, or are omitted altogether, thus missing an opportunity to clarify some mysterious points. This includes Vogel’s construction of a non-Lie-algebra weight system and the Goussarov–Polyak–Viro proof of the existence of “Gauss diagram formulas”.

**What I wish there was in the book but there isn’t.** The relationship with Chern–Simons theory, Feynman diagrams, and configuration space integrals, culminating in an alternative (and more “3D”) proof of the Fundamental Theorem. This is a major omission.

**Why I hope there will be a continuation book one day.** There’s much more to the story! There are finite type invariants of 3-manifolds, and of certain classes of 2-dimensional knots in  $\mathbb{R}^4$ , and of “virtual knots”, and they each have their lovely yet non-obvious theories, and these theories link with each other and with other branches of Lie theory, algebra, topology, and quantum field theory. Volume 2 is sorely needed.



REFERENCES

- [1] Dror Bar-Natan, *On the Vassiliev knot invariants*, *Topology* **34** (1995), no. 2, 423–472, DOI 10.1016/0040-9383(95)93237-2. MR1318886 (97d:57004)
- [2] D. Bar-Natan, *Finite Type Invariants*, in *Encyclopedia of Mathematical Physics*, (J.-P. Francoise, G. L. Naber and Tsou S. T., eds.) Elsevier, Oxford, 2006 (vol. 2 p. 340).
- [3] S. Chmutov, S. Duzhin, and J. Mostovoy, *Introduction to Vassiliev knot invariants*, Cambridge University Press, Cambridge, 2012. MR2962302
- [4] Joan S. Birman and Xiao-Song Lin, *Knot polynomials and Vassiliev's invariants*, *Invent. Math.* **111** (1993), no. 2, 225–270, DOI 10.1007/BF01231287. MR1198809 (94d:57010)
- [5] Predrag Cvitanović, *Group theory*, Princeton University Press, Princeton, NJ, 2008. Birdtracks, Lie's, and exceptional groups. MR2418111 (2010b:22001)
- [6] M. N. Gusarov, *A new form of the Conway-Jones polynomial of oriented links*, *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* **193** (1991), no. Geom. i Topol. 1, 4–9, 161 (Russian, with English summary). MR1157140 (93b:57007)
- [7] M. Gusarov, *On  $n$ -equivalence of knots and invariants of finite degree*, *Topology of manifolds and varieties*, *Adv. Soviet Math.*, vol. 18, Amer. Math. Soc., Providence, RI, 1994, pp. 173–192. MR1296895 (96i:57005)
- [8] Vaughan F. R. Jones, *A polynomial invariant for knots via von Neumann algebras*, *Bull. Amer. Math. Soc. (N.S.)* **12** (1985), no. 1, 103–111, DOI 10.1090/S0273-0979-1985-15304-2. MR766964 (86e:57006)
- [9] Maxim Kontsevich, *Vassiliev's knot invariants*, *I. M. Gel'fand Seminar*, *Adv. Soviet Math.*, vol. 16, Amer. Math. Soc., Providence, RI, 1993, pp. 137–150. MR1237836 (94k:57014)
- [10] Maxim Kontsevich, *Feynman diagrams and low-dimensional topology*, *First European Congress of Mathematics, Vol. II (Paris, 1992)*, *Progr. Math.*, vol. 120, Birkhäuser, Basel, 1994, pp. 97–121. MR1341841 (96h:57027)
- [11] Roger Penrose, *Applications of negative dimensional tensors*, *Combinatorial Mathematics and its Applications (Proc. Conf., Oxford, 1969)*, Academic Press, London, 1971, pp. 221–244. MR0281657 (43 #7372)
- [12] N. Yu. Reshetikhin and V. G. Turaev, *Ribbon graphs and their invariants derived from quantum groups*, *Comm. Math. Phys.* **127** (1990), no. 1, 1–26. MR1036112 (91c:57016)
- [13] V. A. Vassiliev, *Cohomology of knot spaces*, *Theory of singularities and its applications*, *Adv. Soviet Math.*, vol. 1, Amer. Math. Soc., Providence, RI, 1990, pp. 23–69. MR1089670 (92a:57016)
- [14] V. A. Vassiliev, *Complements of discriminants of smooth maps: topology and applications*, *Translations of Mathematical Monographs*, vol. 98, American Mathematical Society, Providence, RI, 1992. Translated from the Russian by B. Goldfarb. MR1168473 (94i:57020)
- [15] Edward Witten, *Quantum field theory and the Jones polynomial*, *Comm. Math. Phys.* **121** (1989), no. 3, 351–399. MR990772 (90h:57009)

DROR BAR-NATAN  
UNIVERSITY OF TORONTO, CANADA

# American Mathematical Society

Job ID: mstp-97

Title: /ams/produ

User: kta


Date: 28 Feb 2013 08:16:35

Author: This scan of the marked-up proofs of your paper is for informational purposes only. Please do not mark corrections on this copy, and do not return this copy. It will not be checked for corrections. All corrections should be marked on the author galleys that are to be returned to Providence.

od/bull1413/bull1413.dvi



**AMS**





BULLETIN (New Series) OF THE  
 AMERICAN MATHEMATICAL SOCIETY  
 S 0273-0979(XX)0000-0

*Introduction to Vassiliev knot invariants*, by S. Chmutov, S. Duzhin, and J. Mostovoy,  
 Cambridge University Press, Cambridge, UK, 2012, xvi+504 pp., ISBN 978-1-107029083-2, hardback, US \$70.00.

Merely 30 years ago, if you had asked even the best informed mathematician about the relationship between knots and Lie algebras, she would have laughed, for there isn't and there can't be. Knots are flexible, Lie algebras are rigid. Knots are irregular, Lie algebras are symmetric. The list of knots is a lengthy mess, the collection of Lie algebras is well-organized. Knots are useful for sailors, scouts, and hangmen, Lie algebras for navigators, engineers, and high energy physicists. Knots are blue collar, Lie algebras are white. They are as similar as worms and crystals: both well-studied, but hardly ever together. Then in the 1980s came Jones, and




FIGURE 1. A knot and a Lie algebra, a list of knots and a list of Lie algebras, and an unusual conference of the symmetric and the knotted.

Witten, and Reshetikhin and Turaev [8, 12, 15] and showed that if you really are the best informed, and you know your quantum field theory and conformal field theory and quantum groups, then you know that the two disjoint fields are in fact intricately related. This “quantum” approach remains the most powerful way to get computable knot invariants out of (certain) Lie algebras (and representations thereof). Yet shortly later, in the late 80s and early 90s, an alternative perspective aroused, that of “finite-type” or “Vassiliev-Goussarov” invariants [1, 4, 6, 7, 9, 10, 13, 14], which made the surprising relationship between knots and Lie algebras appear simple and almost inevitable.

The reviewed [3] is about that alternative perspective, the one reasonable sounding but not entirely trivial theorem that is crucially needed within it (the “Fundamental Theorem” or the “Kontsevich integral”), and the many threads that begin

2010 *Mathematics Subject Classification*. Primary 57M25.

Picture credits: Rope from “The Project Gutenberg eBook, Knots, Splices and Rope Work, by A. Hyatt Verrill”, <http://www.gutenberg.org/files/13510/13510-h/13510-h.htm>. Plane from NASA, <http://www.grc.nasa.gov/WWW/k-12/airplane/rotations.html>.

TeX at <http://drorbn.net/AcademicPensieve/2013-01/CDMReview/>, copleft at <http://www.math.toronto.edu/~drorbn/Copleft/>. This review was written while I was a guest at the Newton Institute, in Cambridge, UK..

blue/hyphen  
 = 1/2 / cover  
 58  
 #  
 #  
 permission?  
 no-persivg  
 first use.

192/11  
 1/11  
 book order / 3  
 3/11/13 (20)

?

changed per CTP

with that perspective. Let me start with a brief summary of the mathematics, and even before, an even briefer summary.

The briefer summary is that in some combinatorial sense it is possible to “differentiate” knot invariants, and hence it makes sense to talk about “polynomials” on the space of knots — these are functions on the set of knots (namely, these are knot invariants) whose sufficiently high derivatives vanish. Such polynomials can be fairly conjectured to separate knots — elsewhere in math in lucky cases polynomials separate points, and in our case, specific computations are encouraging. Also, such polynomials are determined by their “coefficients”, and each of these, by the one-side-easy “Fundamental Theorem”, is a linear functional on some finite space of graphs modulo relations. These same graphs turn out to parameterize formulas that make sense in a wide class of Lie algebras, and the said relations match exactly with the relations in the definition of a Lie algebra — anti-symmetry and the Jacobi identity. Hence what is more or less dual to knots (invariants), is also, after passing to the coefficients, dual to certain graphs which are more or less dual to Lie algebras. QED, and on to the less brief summary.

Let  $V$  be an arbitrary invariant of oriented knots in oriented space with values in (say)  $\mathbb{Q}$ . Extend  $V$  to be an invariant of 1-singular knots, knots that have a single singularity that locally looks like a double point  $\times$ , using the formula

$$(1) \quad V(\times) = V(\curvearrowright) - V(\curvearrowleft).$$

Further extend  $V$  to the set  $\mathcal{K}^m$  of  $m$ -singular knots (knots with  $m$  such double points) by repeatedly using (1).

**Definition 1.** We say that  $V$  is of type  $m$  (or “Vassiliev of type  $m$ ”) if its extension  $V|_{\mathcal{K}^{m+1}}$  to  $(m+1)$ -singular knots vanishes identically. We say that  $V$  is of finite type (or “Vassiliev”) if it is of type  $m$  for some  $m$ .

Repeated differences are similar to repeated derivatives and hence it is fair to think of the definition of  $V|_{\mathcal{K}^m}$  as repeated differentiation. With this in mind, the above definition imitates the definition of polynomials of degree  $m$ . Hence finite type invariants can be thought of as “polynomials” on the space of knots. It is known (see e.g. [3]) that the class of finite type invariants is large and powerful. Yet the first question on finite type invariants remains unanswered:

**Problem 2.** Honest polynomials are dense in the space of functions. Are finite type invariants dense within the space of all knot invariants? Do they separate knots?

The top derivatives of a multi-variable polynomial form a system of constants that determine that polynomial up to polynomials of lower degree. Likewise the  $m$ th derivative  $V^{(m)} = V|_{\mathcal{K}^m} = V(\times^{\cdot m} \times)$  of a type  $m$  invariant  $V$  is a constant in the sense that it does not see the difference between overcrossings and undercrossings and so it is blind to 3D topology. Indeed  $V(\times^{\cdot m} \times \times) - V(\times^{\cdot m} \times \times) =$

<sup>1</sup>Partially self-plagiarized from [2].

<sup>2</sup>Keep this apart from invariants of knots whose values are polynomials, such as the Alexander or the Jones polynomial. A posteriori related, these are a priori entirely different.

<sup>3</sup>As common in the knot theory literature, in the formulas that follow a picture such as  $\times^{\cdot m} \times$  indicates “some knot having  $m$  double points and a further (right-handed) crossing”. Furthermore, when two such pictures appear within the same formula, it is to be understood that the parts of the knots (or diagrams) involved *outside* of the displayed pictures are to be taken as the same.

combinatorics

5/11

(A)

(A)

(A)

(A)

7

(A)

5/11

do not put across page







Figure 1

see orig. ms for figure placement


$V(\bigotimes^{m+1} X) = 0$ . Also, clearly  $V^{(m)}$  determines  $V$  up to invariants of lower type. Hence a primary tool in the study of finite type invariants is the study of the “top derivative”  $V^{(m)}$ , also known as “the weight system of  $V$ ”. Blind to 3D topology,  $V^{(m)}$  only sees the combinatorics of the circle that parameterizes an  $m$ -singular knot. On this circle there are  $m$  pairs of points that are pairwise identified in the image; standardly one indicates those by drawing a circle with  $m$  chords marked (an “ $m$ -chord diagram”) as on the right. Let  $\mathcal{D}_m$  denote the space of all formal linear combinations with rational coefficients of  $m$ -chord diagrams. Thus  $V^{(m)}$  is a linear functional on  $\mathcal{D}_m$ .

I leave it for the reader to figure out or read in [3, pp. 88] how the following figure easily implies the “4T” relations of the “easy side” of the theorem that follows:



**Theorem 3.** (The Fundamental Theorem, details in [3]).

- (Easy side) If  $V$  is a rational valued type  $m$  invariant then  $V^{(m)}$  satisfies the “4T” relations shown on the right, and hence it descends to a linear functional on  $\mathcal{A}_m := \mathcal{D}_m/4T$ .  
If in addition  $V^{(m)} \equiv 0$ , then  $V$  is of type  $m - 1$ .



- (Hard side, slightly misstated by avoiding “framings”) For any linear functional  $W$  on  $\mathcal{A}_m$  there is a rational valued type  $m$  invariant  $V$  so that  $V^{(m)} = W$ .

Thus to a large extent the study of finite type invariants is reduced to the finite (though super-exponential in  $m$ ) algebraic study of  $\mathcal{A}_m$ .

Much of the richness of finite type invariants stems from their relationship with Lie algebras. Theorem 4 below suggests this relationship on an abstract level and Theorem 5 makes that relationship concrete.

**Theorem 4.** [1] The space  $\mathcal{A}_m$  is isomorphic to the space  $\mathcal{A}_m^t$  generated by “Jacobi diagrams in a circle” (chord diagrams that are also allowed to have oriented internal trivalent vertices) that have exactly  $2m$  vertices, modulo the AS, STU and IHX relations. See the figure on the right.

The key to the proof of Theorem 4 is the figure on the right, which shows that the 4T relation is a consequence of two STU relations. The rest is more or less an exercise in induction.



Thinking of internal trivalent vertices as graphical analogs of the Lie bracket, the AS relation becomes the anti-commutativity of the bracket, STU becomes the equation  $[x, y] = xy - yx$  and IHX becomes the Jacobi identity. This analogy

Tr. under


placemat  
AU: de?

?

3 //




A Jacobi diagram in a circle



is made concrete within the following construction, originally due to Penrose [11] and to Cvitanović [5]. Given a finite dimensional metrized Lie algebra  $\mathfrak{g}$  (e.g., any semi-simple Lie algebra) and a finite-dimension representation  $\rho : \mathfrak{g} \rightarrow \text{End}(V)$  of  $\mathfrak{g}$ , choose an orthonormal basis  $\{X_a\}_{a=1}^{\dim \mathfrak{g}}$  of  $\mathfrak{g}$  and some basis  $\{v_\alpha\}_{\alpha=1}^{\dim V}$  of  $V$ , let  $f_{abc}$  and  $r_{\alpha\beta}^\gamma$  be the “structure constants” defined by

$$f_{abc} := \langle [X_a, X_b], X_c \rangle \quad \text{and} \quad \rho(X_a)(v_\beta) = \sum_\gamma r_{\alpha\beta}^\gamma v_\gamma.$$

Now given a Jacobi diagram  $D$  label its circle-arcs with Greek letters  $\alpha, \beta, \dots$ , and its chords with Latin letters  $a, b, \dots$ , and map it to a sum as suggested by the following example:



**Theorem 5.** *This construction is well defined, and the basic properties of Lie algebras imply that it respects the AS, STU, and IHX relations. Therefore it defines a linear functional  $W_{\mathfrak{g},R} : \mathcal{A}_m \rightarrow \mathbb{Q}$ , for any  $m$ .*

The last assertion along with Theorem 3 show that associated with any  $\mathfrak{g}$ ,  $R$  and  $m$  there is a weight system and hence a knot invariant. Thus knots are indeed linked with Lie algebras.

The above is of course merely a sketch of the beginning of a long story. You can read the details, and some of the rest, in [3].

# **What I like about [3].** Detailed, well thought out, and carefully written. Lots of pictures! Many excellent exercises! A complete discussion of “the algebra of chord diagrams”. A nice discussion of the pairing of diagrams with Lie algebras, including examples aplenty. The discussion of the Kontsevich integral (meaning, the proof of the hard side of Theorem 3) is terrific — detailed and complete and full

<sup>4</sup>This requirement can easily be relaxed.

(P)

Proposition

Counter



of pictures and examples, adding a great deal to the original sources. The subject of “associators” is huge and worthy of its own book(s); yet in as much as they are related to Vassiliev invariants, the discussion in [3] is excellent. A great many further topics are touched — multiple  $\zeta$ -values, the relationship of the Hopf link with the Duflo isomorphism, intersection graphs and other combinatorial aspects of chord diagrams, Rozansky’s rationality conjecture, the Melvin-Morton conjecture, braids,  $n$ -equivalence, etc.

For all these, I’d certainly recommend [3] to any newcomer to the subject of knot theory, starting with my own students.

However, some proofs other than that of Theorem 3 are repeated as they appear in original articles with only a superficial touch-up, or are omitted altogether, thus missing an opportunity to clarify some mysterious points. This includes Vogel’s construction of a non-Lie-algebra weight system and the Goussarov-Polyak-Viro proof of the existence of “Gauss diagram formulas”.

# What I wish there was in the book, but there isn’t. The relationship with Chern-Simons theory, Feynman diagrams, and configuration space integrals, culminating in an alternative (and more “3D”) proof of the Fundamental Theorem. This is a major omission.

# Why I hope there will be a continuation book, one day. There’s much more to the story! There are finite type invariants of 3-manifolds, and of certain classes of 2-dimensional knots in  $\mathbb{R}^4$ , and of “virtual knots”, and they each have their lovely yet non-obvious theories, and these theories link with each other and with other branches of Lie theory, algebra, topology, and quantum field theory. Volume 2 is sorely needed.

## REFERENCES

- [1] Dror Bar-Natan, *On the Vassiliev knot invariants*, *Topology* **34** (1995), no. 2, 423–472, DOI 10.1016/0040-9383(95)93237-2. MR1318886 (97d:57004)
- [2] D. Bar-Natan, *Finite Type Invariants*, in *Encyclopedia of Mathematical Physics*, (J.-P. Francoise, G. L. Naber and Tsou S. T., eds.) Elsevier, Oxford, 2006 (vol. 2 p. 340).
- [3] The reviewed book.
- [4] Joan S. Birman and Xiao-Song Lin, *Knot polynomials and Vassiliev’s invariants*, *Invent. Math.* **111** (1993), no. 2, 225–270, DOI 10.1007/BF01231287. MR1198809 (94d:57010)
- [5] Predrag Cvitanović, *Group theory*, Princeton University Press, Princeton, NJ, 2008. Birdtracks, Lie’s, and exceptional groups. MR2418111 (2010b:22001)
- [6] M. N. Gusarov, *A new form of the Conway-Jones polynomial of oriented links*, *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* **193** (1991), no. Geom. i Topol. 1, 4–9, 161 (Russian, with English summary). MR1157140 (93b:57007)
- [7] M. Gusarov, *On  $n$ -equivalence of knots and invariants of finite degree*, *Topology of manifolds and varieties*, *Adv. Soviet Math.*, vol. 18, Amer. Math. Soc., Providence, RI, 1994, pp. 173–192. MR1296895 (96i:57005)
- [8] Vaughan F. R. Jones, *A polynomial invariant for knots via von Neumann algebras*, *Bull. Amer. Math. Soc. (N.S.)* **12** (1985), no. 1, 103–111, DOI 10.1090/S0273-0979-1985-15304-2. MR766964 (86e:57006)
- [9] Maxim Kontsevich, *Vassiliev’s knot invariants*, *I. M. Gel’fand Seminar*, *Adv. Soviet Math.*, vol. 16, Amer. Math. Soc., Providence, RI, 1993, pp. 137–150. MR1237836 (94k:57014)
- [10] Maxim Kontsevich, *Feynman diagrams and low-dimensional topology*, *First European Congress of Mathematics*, Vol. II (Paris, 1992), *Progr. Math.*, vol. 120, Birkhäuser, Basel, 1994, pp. 97–121. MR1341841 (96h:57027)
- [11] Roger Penrose, *Applications of negative dimensional tensors*, *Combinatorial Mathematics and its Applications* (Proc. Conf., Oxford, 1969), Academic Press, London, 1971, pp. 221–244. MR0281657 (43 #7372)

S. Chmutov, S. Duval, and J. Mostrovo, *Introduction to Vassiliev knot invariants*, Cambridge University Press, Cambridge, 2012. MR 2962302.

- [12] N. Yu. Reshetikhin and V. G. Turaev, *Ribbon graphs and their invariants derived from quantum groups*, *Comm. Math. Phys.* **127** (1990), no. 1, 1–26. MR1036112 (91c:57016)
- [13] V. A. Vassiliev, *Cohomology of knot spaces*, *Theory of singularities and its applications*, *Adv. Soviet Math.*, vol. 1, Amer. Math. Soc., Providence, RI, 1990, pp. 23–69. MR1089670 (92a:57016)
- [14] V. A. Vassiliev, *Complements of discriminants of smooth maps: topology and applications*, *Translations of Mathematical Monographs*, vol. 98, American Mathematical Society, Providence, RI, 1992. Translated from the Russian by B. Goldfarb. MR1168473 (94i:57020)
- [15] Edward Witten, *Quantum field theory and the Jones polynomial*, *Comm. Math. Phys.* **121** (1989), no. 3, 351–399. MR990772 (90h:57009)

DOR BAR-NATAN  
UNIVERSITY OF TORONTO, CANADA