


Dror Bar-Natan: Talks: Bern-131104:

$\omega := \text{http://www.math.toronto.edu/~drorbn/Talks/Bern-131104}$
Video, handout, links at $\omega /$



The Kashiwara-Vergne Problem and Topology

Abstract. I will describe a general machine, a close cousin of Taylor's theorem, whose inputs are topics in topology and whose outputs are problems in algebra. There are many inputs the machine can take, and many outputs it produces, but I will concentrate on just one input/output pair. When fed with a certain class of knotted 2-dimensional objects in 4-dimensional space, it outputs the Kashiwara-Vergne Problem (1978 ω/KV , solved Alekseev-Meinrenken 2006 ω/AM , elucidated Alekseev-Torossian 2008-2012 ω/AT), a problem about convolutions on Lie groups and Lie algebras.



The Kashiwara-Vergne Conjecture. There exist two series F and G in the completed free Lie algebra FL in generators x and y so that $x+y-\log e^y e^x = (1-e^{-\text{ad } x})F + (e^{\text{ad } y}-1)G$ in FL and so that with $z = \log e^x e^y$,

$$\text{tr}(\text{ad } x)\partial_x F + \text{tr}(\text{ad } y)\partial_y G \text{ in cyclic words} \\ = \frac{1}{2} \text{tr} \left(\frac{\text{ad } x}{e^{\text{ad } x}-1} + \frac{\text{ad } y}{e^{\text{ad } y}-1} - \frac{\text{ad } z}{e^{\text{ad } z}-1} - 1 \right)$$

Implies the loosely-stated **convolutions statement**: Convolutions of invariant functions on a Lie group agree with convolutions of invariant functions on its Lie algebra.


The Machine. Let G be a group, $\mathcal{K} = \mathbb{Q}G = \{\sum a_i g_i : a_i \in \mathbb{Q}, g_i \in G\}$ its group-ring, $\mathcal{I} = \{\sum a_i g_i : \sum a_i = 0\} \subset \mathcal{K}$ its augmentation ideal. Let

$$\mathcal{A} = \text{gr } \mathcal{K} := \bigoplus_{m \geq 0} \mathcal{I}^m / \mathcal{I}^{m+1}.$$

Note that \mathcal{A} inherits a product from G .
Definition. A linear $Z: \mathcal{K} \rightarrow \mathcal{A}$ is an "expansion" if for any $\gamma \in \mathcal{I}^m$, $Z(\gamma) = (0, \dots, 0, \gamma / \mathcal{I}^{m+1}, *, \dots)$, and a "homomorphic expansion" if in addition it preserves the product.




Example. Let $\mathcal{K} = C^\infty(\mathbb{R}^n)$ and $\mathcal{I} = \{f: f(0) = 0\}$. Then $\mathcal{I}^m = \{f: f \text{ vanishes like } |x|^m\}$ so $\mathcal{I}^m / \mathcal{I}^{m+1}$ degree m homogeneous polynomials and $\mathcal{A} = \{\text{power series}\}$. The Taylor series is a homomorphic expansion!

Just for fun. $\mathcal{K} = \left\{ \begin{matrix} \text{img of bear} \\ \omega/\text{Bear} \end{matrix} \right\} = \left(\text{The set of all 2D projections of reality} \right)$



In the finitely presented case, finding Z amounts to solving a system of equations in a graded space.

Theorem (with Zsuzsanna Dancso, ω/WKO). There is a bijection between the set of homomorphic expansions for $w\mathcal{K}$ and the set of solutions of the Kashiwara-Vergne problem. **This is the tip of a major iceberg.**



The Machine generalizes to arbitrary algebraic structures!
 ω/mac
"God created the knots, all else in topology is the work of mortals."
Leopold Kronecker (modified) www.katlas.org The Knot-Films

colour correction
down correction

A
no, due to lack of space.

A ... major iceberg: associators (8) and elliptic
associators (7), Grothendieck-Taichmüller groups (9),
quantization of Lie bialgebras (10),
deformation quantization of Poisson structures (11), ...