

Video and more @ Dror Bar-Natan: Talks: Vienna-1402:  
<http://www.math.toronto.edu/~drorbn/Talks/Vienna-1402>

**A Partial Reduction of BF Theory to Combinatorics, I**

**Abstract.** I will describe a semi-rigorous reduction to computable combinatorics of perturbative BF theory (Cattaneo-Rossi [CR]) in the case of ribbon 2-links. Also, I will explain how and why my approach may or may not work in the non-ribbon case. Weak this result is, and at least partially already known (Watanabe [Wa]). Yet in the ribbon case, the resulting invariant is a universal finite type invariant, a gadget that significantly generalizes and clarifies the Alexander polynomial and that is closely related to the Kashiwara-Vergne problem. I cannot rule out the possibility that the corresponding gadget in the non-ribbon case will be as interesting. *(good news in highlight)*

**The BF Feynman Rules.** For an edge  $e$ , let  $\Phi_e$  be its direction, in  $S^3$  or  $S^1$ . Let  $\omega_3$  and  $\omega_1$  be volume forms on  $S^3$  and  $S^1$ . Then for a 2-link  $(f_i)_{i \in T}$ ,



$$\zeta = \log \sum_{\text{diagrams } D} \frac{D}{|\text{Aut}(D)|} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^4} \int_{\mathbb{R}^4} \prod_{\text{red } e \in D} \Phi_e^* \omega_3 \prod_{\text{black } e \in D} \Phi_e^* \omega_1$$

is an invariant in  $CW(FL(T)) \rightarrow CW(T)$ , "cyclic words in  $T$ ".


**BF Following [CR].**  $A \in \Omega^1(M = \mathbb{R}^4, \mathfrak{g})$ ,  $B \in \Omega^2(M, \mathfrak{g}^*)$ ,

$$S(A, B) := \int_M \langle B, F_A \rangle.$$


With  $f: (S = \mathbb{R}^2) \rightarrow M$ ,  $\xi \in \Omega^0(S, \mathfrak{g})$ ,  $\beta \in \Omega^1(S, \mathfrak{g}^*)$ , set

$$O(A, B, f) := \int \mathcal{D}\xi \mathcal{D}\beta \exp\left(\frac{i}{\hbar} \int_S \langle \xi, d_f A + f^* B \rangle\right).$$


**Decker Sets ("2D Gauss Codes").**





**Some Examples.**



**A BF Feynman Diagram.**



*Go over invariance proof.*



## A Partial Reduction of BF Theory to Combinatorics, 2

**Theorem 1 (with Cattaneo (credit, no blame)).** In the ribbon case,  $e^c$  can be computed as follows:

$$e^c \leftarrow \sum_{k,l,m \geq 0} \frac{(+)^k (-)^l (+)^m}{k!l!m!}$$


$$e^c \leftarrow \sum_{k,m \geq 0} \frac{(+)^k (-)^m}{k!m!}$$

**Theorem 2.** Using Gauss diagrams to represent knots and  $T$ -component pure tangles, the above formulas define an invariant in  $CW(FL(T)) \rightarrow CW(T)$ , "cyclic words in  $T$ ".


- Agrees with BN-Dancso [BND] and with [BN1].
- In-practice computable!
- Vanishes on braids.
- Extends to w.
- Contains Alexander.
- The "missing factor" in Levine's factorization [Le] (the rest of [Le] also fits, hence contains the MVA).
- Related to / extends Farber's [Fa]?
- Should be summed and categorized.

### References.

[Ar] V. I. Arnold, *Topological Invariants of Plane Curves and Caustics*, University Lecture Series 5, American Mathematical Society 1994.  
 [BN1] D. Bar-Natan, *Balloons and Hoops and their Universal Finite Type Invariant, BF Theory, and an Ultimate Alexander Invariant*, <http://www.math.toronto.edu/~drorbn/papers/KBH/>, arXiv:1308.1721.  
 [BN2] D. Bar-Natan, *Bracelets and the Goussarov filtration of the space of knots, Invariants of knots and 3-manifolds (Kyoto 2001)*, Geometry and Topology Monographs 4 1-12, arXiv:math.GT/0111267.  
 [BND] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects: From Alexander to Kashiwara and Vergne*, <http://www.math.toronto.edu/~drorbn/papers/WK0/>.  
 [CKS] J. S. Carter, S. Kamada, and M. Saito, *Diagrammatic Computations for Quandles and Cocycle Knot Invariants*, *Contemp. Math.* 318 (2003) 51-74.  
 [CS] J. S. Carter and M. Saito, *Knotted surfaces and their diagrams*, *Mathematical Surveys and Monographs* 55, American Mathematical Society, Providence 1998.  
 [Da] E. Dalvit, <http://science.unitn.it/~dalvit/>.  
 [CR] A. S. Cattaneo and C. A. Rossi, *Wilson Surfaces and Higher Dimensional Knot Invariants*, *Commun. in Math. Phys.* 256-3 (2005) 513-537, arXiv:math-ph/0210037.  
 [Fa] M. Farber, *Noncommutative Rational Functions and Boundary Links*, *Math. Ann.* 293 (1992) 543-568.  
 [Le] J. Levine, *A Factorization of the Conway Polynomial*, *Comment. Math. Helv.* 74 (1999) 27-53, arXiv:q-alg/9711007.  
 [Ro] D. Roseman, *Reidemeister-Type Moves for Surfaces in Four-Dimensional Space*, *Knot Theory*, Banach Center Publications 42 (1998) 347-380.  
 [Wa] T. Watanabe, *Configuration Space Integrals for Long n-Knots, the Alexander Polynomial and Knot Space Cohomology*, *Alg. and Geom. Top.* 7 (2007) 47-92, arXiv:math/0609742.




**Sketch of Proof.** In 4D axial gauge, only "drop down" red propagators, hence in the ribbon case, no  $M$ -trivalent vertices.  $S$  integrals are  $\pm 1$  iff "ground pieces" run on nested curves as below, and exponentials arise when several propagators compete for the same double curve. And then the combinatorics is obvious...




### Musings

**Chern-Simons.** When the domain of BF is restricted to ribbon knots, and the target of CS is restricted to trees and wheels, they agree. Why?


Is this all? What about the  $v$ -invariant? (the "true" triple linking number)



**Gnots.** In 3D, a generic immersion of  $S^1$  is an embedding, a knot. In 4D, a generic immersion of a surface has finitely-many double points (a gnot?). Perhaps we should be studying these?




**Finite type.** What are finite-type invariants for 2-knots? What would be "chord diagrams"?



**Bubble-wrap-finite-type.**

There's an alternative definition of finite type in 3D, due to Goussarov (see [BN2]). The obvious parallel in 4D involves "bubble wraps". Is it any good?



**Shielded tangles.** In 3D, one can't zoom in and compute "the Chern-Simons invariant of a tangle". Yet there are well-defined invariants of "shielded tangles", and rules for their compositions. What would the 4D analog be?



Will the relationship with the Kashiwara-Vergne problem [BND] necessarily arise here?

**Plane curves.** Shouldn't we understand integral / finite type invariants of plane curves, in the style of Arnold's  $J^+$ ,  $J^-$ , and  $St$  [Ar], a bit better?

	$a(\times)$	$a(\times)$	$a(\times)$	$\infty$	$\circ$	$\circ$	$\circ$	$\circ$	$\dots$
$St$	1	0	0	0	0	1	2	3	$\dots$
$J^+$	0	2	0	0	0	-2	-4	-6	$\dots$
$J^-$	0	0	-2	-1	0	-3	-6	-9	$\dots$

"God created the knots, all else in topology is the work of mortals."  
 Leopold Kronecker (modified) [www.katlas.org](http://www.katlas.org)

~~Lego picture?~~

virtual 2-knots? Group only?  
 Find a way to insert YouTube's of flying vortex rings & of knotted vortices? Maybe in place of "propaganda"?

- <https://www.youtube.com/watch?v=mHyTOcfF99o#t=3m45s>
- <http://youtu.be/mHyTOcfF99o#t=3m45s>
- <http://www.youtube.com/watch?v=YCAOVIEVhg&feature=youtu.be#t=50s>
- <http://youtu.be/YCAOVIEVhg#t=50s>

Improve logging at HardantBrowser.js? ✓