

cod-2 cod-2!

perhaps, "is there a finite-type codimension 2 knot theory?"

A Partial Reduction of BF Theory to Combinatorics, I

Abstract. I will describe a semi-rigorous reduction of perturbative BF theory (Cattaneo-Rossi [CR]) to computable combinatorics, in the case of ribbon 2-links. Also, I will explain how and why my approach may or may not work in the non-ribbon case. **Weak** this result is, and at least partially already known (Watanabe [Wa]). Yet in the ribbon case, the resulting invariant is a universal finite type invariant, a gadget that significantly generalizes and clarifies the Alexander polynomial and that is closely related to the Kashiwara-Vergne problem. I cannot rule out the possibility that the corresponding gadget in the non-ribbon case will be as interesting. (good news in highlight)

The BF Feynman Rules. For an edge e , let Φ_e be its direction, in S^1 or S^3 . Let ω_{e_1} and ω_{e_2} be volume forms on S^1 and S^1 . Then for a 2-link $(\kappa_i)_{i \in T}$,

$$\zeta = \log \sum_{\text{diagrams } D} \frac{|D|}{|\text{Aut}(D)|} \int_{\mathbb{R}^2} \dots \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \dots \int_{\mathbb{R}^2} \prod_{\text{red } e \in D} \Phi_e^* \omega_{e_1} \prod_{\text{black } e \in D} \Phi_e^* \omega_{e_2}$$

is an invariant in $CW(FL(T)) \rightarrow CW(T)/\sim$, "symmetrized cyclic words in T ".

BF Following [CR]. $A \in \Omega^1(M = \mathbb{R}^d, \mathfrak{g})$, $B \in \Omega^2(M, \mathfrak{g}^*)$,

$$S(A, B) := \int \langle B, F_A \rangle.$$

With $\kappa: (S = \mathbb{R}^2) \rightarrow M$, $\beta, \alpha \in \Omega^1(S, \mathfrak{g})$, $\alpha \in \Omega^1(S, \mathfrak{g}^*)$, set

$$O(A, B, \kappa) := \int \mathcal{D}\beta \mathcal{D}\alpha \exp\left(\frac{i}{\hbar} \int_S \langle \beta, d_{e_A} \alpha + \kappa^* B \rangle\right).$$

Decker Sets ("2D Gauss Codes").

(only double curves are allowed in ribbon 2-knots)

"a double curve"

"a triple point"

"a branch point"

Carter Banach Saito

Some Examples.

A 4D knot by Carter and Saito [CS]

A 4D knot by Dalvit [Da]

"a w-knot"

A 2-link

"ribbed cigar presentation"

A 2-twist spun trefoil by Carter-Kamada-Saito [CKS].

A BF Feynman Diagram.

split to a link
remove all rattles.

trees on wheels, odd edges even rattles

degree = #(rattles)

air piece
rattle
snake

Ro1
Ro2
Ro3
Ro4
Ro5
Ro6
Ro7

Roseman [Ro]



Cattaneo Rossi

PHOTO NOT AVAILABLE


QR code

A Partial Reduction of BF Theory to Combinatorics, 2

Theorem 1 (with Cattaneo, Dalvit (credit, no blame)). In the ribbon case, $e^{\mathcal{L}}$ can be computed as follows:




Sketch of Proof. In 4D axial gauge, only "drop down" red propagators, hence in the ribbon case, no M -trivalent vertices. S integrals are ± 1 iff "ground pieces" run on nested curves as below, and exponentials arise when several propagators compete for the same double curve. And then the combinatorics is obvious...



Musings

Chern-Simons. When the domain of BF is restricted to ribbon knots, and the target of Chern-Simons is restricted to trees and wheels, they agree. Why?

Is this all? What about the ν -invariant?




Gnots. In 3D, a generic immersion of S^1 is an embedding, a knot. In 4D, a generic immersion of a surface has finitely-many double points (a gnot?). Perhaps we should be studying these?

Finite type. What are finite-type invariants for 2-knots? What would be "chord diagrams"?



Bubble-wrap-finite-type.

There's an alternative definition of finite type in 3D, due to Goussarov (see [BN1]). The obvious parallel in 4D involves "bubble wraps". Is it any good?



Shielded tangles. In 3D, one can't zoom in and compute "the Chern-Simons invariant of a tangle". Yet there are well-defined invariants of "shielded tangles", and rules for their compositions. What would the 4D analog be?



Will the relationship with the Kashiwara-Vergne problem [BND] necessarily arise here?

Plane curves. Shouldn't we understand integral / finite type invariants of plane curves, in the style of Arnold's J^+ , J^- , and St [Ar], a bit better?



| | $a(\times)$ | $a(\times)$ | $a(\times)$ | ∞ | \circ | \circ | \circ | \circ | \circ | \dots |
|-------|-------------|-------------|-------------|----------|---------|---------|---------|---------|---------|---------|
| St | 1 | 0 | 0 | 0 | 0 | 1 | 2 | 3 | \dots | \dots |
| J^+ | 0 | 2 | 0 | 0 | 0 | -2 | -4 | -6 | \dots | \dots |
| J^- | 0 | 0 | -2 | -1 | 0 | -3 | -6 | -9 | \dots | \dots |


Theorem 2. Using Gauss diagrams to represent knots and T -component pure tangles, the above formulas define an invariant in $CW(FL(T)) \rightarrow CW(T)$, "cyclic words in T ".

- Agrees with BN-Dancso [BND] and with [BN2].
- In-practice computable!
- Vanishes on braids.
- Extends to w .
- Contains Alexander.
- The "missing factor" in Levine's factorization [Le] (the rest of [Le] also fits, hence contains the MVA).
- Related to / extends Farber's [Fa]?
- Should be summed and categorized.

References

[Ar] V. I. Arnold, *Topological Invariants of Plane Curves and Caustics*, University Lecture Series 5, American Mathematical Society 1994.
 [BN1] D. Bar-Natan, *Bracelets and the Goussarov filtration of the space of knots*, *Invariants of knots and 3-manifolds (Kyoto 2001)*, Geometry and Topology Monographs 4 1–12, arXiv:math.GT/0111267.
 [BN2] D. Bar-Natan, *Balloons and Hoops and their Universal Finite Type Invariant*, *BF Theory, and an Ultimate Alexander Invariant*, <http://www.math.toronto.edu/~drorbn/papers/KBH/>, arXiv:1308.1721.
 [BND] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W -Knotted Objects: From Alexander to Kashiwara and Vergne*, <http://www.math.toronto.edu/~drorbn/papers/WKO/>.
 [CKS] J. S. Carter, S. Kamada, and M. Saito, *Diagrammatic Computations for Quandles and Cocycle Knot Invariants*, *Contemp. Math.* 318 (2003) 51–74.
 [CS] J. S. Carter and M. Saito, *Knotted surfaces and their diagrams*, *Mathematical Surveys and Monographs* 55, American Mathematical Society, Providence 1998.
 [Da] E. Dalvit, <http://science.unitn.it/~dalvit/>.
 [CR] A. S. Cattaneo and C. A. Rossi, *Wilson Surfaces and Higher Dimensional Knot Invariants*, *Commun. in Math. Phys.* 256-3 (2005) 513–537, arXiv:math-ph/0210037.
 [Fa] M. Farber, *Noncommutative Rational Functions and Boundary Links*, *Math. Ann.* 293 (1992) 543–568.
 [Le] J. Levine, *A Factorization of the Conway Polynomial*, *Comment. Math. Helv.* 74 (1999) 27–53, arXiv:q-alg/9711007.
 [Ro] D. Roseman, *Reidemeister-Type Moves for Surfaces in Four-Dimensional Space*, *Knot Theory*, Banach Center Publications 42 (1998) 347–380.
 [Wa] T. Watanabe, *Configuration Space Integrals for Long n -Knots, the Alexander Polynomial and Knot Space Cohomology*, *Alg. and Geom. Top.* 7 (2007) 47–92, arXiv:math/0609742.

Continuing Joost Slingerland...



"God created the knots, all else in topology is the work of mortals."
 Leopold Kronecker (modified)



www.katlas.org

what's going on?