


Abstract. I will describe some very good formulas for a (matrix plus scalar)-valued extension of the Alexander polynomial to tangles, then say that everything extends to virtual tangles, then roughly to simply knotted balloons and hoops in 4D, then the target space extends to (free Lie algebras plus cyclic words), and the result is a universal finite type of the knotted objects in its domain. Taking a cue from the BF topological quantum field theory, everything should extend (with some modifications) to arbitrary codimension-2 knots in arbitrary dimension and in particular, to arbitrary 2-knots in 4D. But what is really going on is still a mystery.

Tangles.




Why Tangles?

- Finitely presented. (meta-associativity: m\_a^{ab} // m\_a^{ac} = m\_b^{bc} // m\_a^{ab})
• Divide and conquer proofs and computations.
• "Algebraic Knot Theory": If K is ribbon,

Z(K) ∈ {cl\_2(Z): cl\_1(Z) = 1}.

(Genus and crossing number are also definable properties).



Theorem 1. ∃! an invariant γ: {pure framed S-component tangles} → R × M\_{S×S}(R), where R = R\_S = ℤ((T\_a)\_{a∈S}) is the ring of rational functions in S variables, intertwining

1. (ω\_1 | S\_1, ω\_2 | S\_2) → (ω\_1 ω\_2 | S\_1 S\_2, S\_1 A\_1 0, S\_2 0 A\_2)

2. (ω | a b S) → (μ ω | c S, c γ + α δ / μ, ε + δ θ / μ, φ + α ψ / μ, Ξ + ψ θ / μ)\_{T\_a, T\_b → T\_c}

and satisfying (|a; a ↗ b, b ↘ a) → (1 | a; a 1 1 - T\_a^{±1}, a | b 0 T\_a^{±1})

In Addition, • This is really "just" a stitching formula for Burau/Gassner [LD, KLW, CT].

- L ↦ ω is Alexander, mod units.
• L ↦ (ω, A) ↦ ω det'(A - I)/(1 - T) is the MVA, mod units.
• The "fastest" Alexander algorithm.
• There are also formulas for strand deletion, reversal, and doubling.
• Every step along the computation is the invariant of something.
• Extends to and more naturally defined on v/w-tangles.
• Fits in one column, including propaganda & implementation.



Implementation key idea:

(ω, A = (α\_{ab})) ↔ (ω, λ = ∑ α\_{ab} t\_a h\_b)
F := F[ω1, λ1] F[ω2, λ2] := F[ω1 ω2, λ1 λ2];
m\_{a,b} := F[ω, λ] := Module[(α, β, γ, δ, θ, ε, φ, ψ, Ξ, μ),
(α β θ | γ δ ε | φ ψ Ξ) = (∂\_{c\_a, h\_a} λ ∂\_{c\_b, h\_b} λ ∂\_{c\_a} λ | ∂\_{c\_b, h\_b} λ ∂\_{c\_a} λ λ) / . (t | h)\_{a1b} → 0;
F[(μ = 1 - β) ω, {t\_c, 1}. (γ + α δ / μ, ε + δ θ / μ, φ + α ψ / μ, Ξ + ψ θ / μ). {h\_a, 1}] / . {T\_a → T\_c, T\_b → T\_c} // FCollect];
FP\_{a,b} := F[1, {t\_a, t\_b}. (1 1 - T\_a | 0 T\_a). {h\_a, h\_b}];
RM\_{a,b} := RP\_{ab} / . T\_a → 1 / T\_a;

Meta-Associativity


γ = F[ω, {t\_1, t\_2, t\_3, t\_s}. (α\_{11} α\_{12} α\_{13} θ\_1 | α\_{21} α\_{22} α\_{23} θ\_2 | α\_{31} α\_{32} α\_{33} θ\_3 | φ\_1 φ\_2 φ\_3 Ξ) . {h\_1, h\_2, h\_3, h\_s}];

(γ // m\_{12→1} // m\_{13→1}) = (γ // m\_{23→2} // m\_{12→1})

True R3


{RM\_{51} RM\_{62} RP\_{34} // m\_{14→1} // m\_{25→2} // m\_{36→3}, RP\_{61} RM\_{24} RM\_{35} // m\_{14→1} // m\_{25→2} // m\_{36→3}}

Matrix equations involving h\_1, h\_2, h\_3 and T\_1, T\_2, T\_3.



γ = RM\_{12,1} RM\_{27} RM\_{83} RM\_{4,11} RP\_{16,5} RP\_{6,13} RP\_{14,9} RP\_{10,15};


Do [γ = γ // m\_{1k→1}, {k, 2, 16}];




Weaknesses, • m\_c^{ab} is non-linear.

- The product ωA is always Laurent, but proving this takes induction with exponentially many conditions.


K^{bh}(H; T).




Examples.



Disturbing Conjecture




Dictionary.




**Operations**

Punctures & Cuts



If  $X$  is a space,  $\pi_1(X)$  is a group,  $\pi_2(X)$  is an Abelian group, and  $\pi_1$  acts on  $\pi_2$ .

**Proposition.** The generators generate.



**Definition.**  $l_{xu}$  is the linking number of hoop  $x$  with balloon  $u$ . For  $x \in H$ ,  $\sigma_x := \prod_{u \in T} T_u^{l_{xu}} \in R = R_T = \mathbb{Z}\langle (T_a)_{a \in T} \rangle$ , the ring of rational functions in  $T$  variables.

**Theorem 2 [BNS].**  $\exists!$  an invariant  $\beta: w\mathcal{K}^{bh}(H; T) \rightarrow R \times M_{T \times H}(R)$ , intertwining

$$1. \left( \begin{array}{c|c} \omega_1 & H_1 \\ \hline T_1 & A_1 \end{array}, \begin{array}{c|c} \omega_2 & H_2 \\ \hline T_2 & A_2 \end{array} \right) \xrightarrow{\sqcup} \begin{array}{c|cc} \omega_1 \omega_2 & H_1 & H_2 \\ \hline T_1 & A_1 & 0 \\ \hline & 0 & A_2 \end{array}$$

$$2. \begin{array}{c|c} \omega & H \\ \hline u & \alpha \\ v & \beta \\ T & \Xi \end{array} \xrightarrow{tm_w^{uv}} \begin{array}{c|c} \omega & H \\ \hline w & \alpha + \beta \\ T & \Xi \end{array}_{T_u, T_v \rightarrow T_w}$$

$$3. \begin{array}{c|cc} \omega & x & y & H \\ \hline T & \alpha & \beta & \Xi \end{array} \xrightarrow{hm_z^{xy}} \begin{array}{c|c} \omega & z & H \\ \hline T & \alpha + \sigma_x \beta & \Xi \end{array}$$

$$4. \begin{array}{c|cc} \omega & x & H \\ \hline u & \alpha & \theta \\ T & \phi & \Xi \end{array} \xrightarrow[\nu:=1+\alpha]{tha^{ux}} \begin{array}{c|cc} \nu\omega & x & H \\ \hline u & \sigma_x \alpha / \nu & \sigma_x \theta / \nu \\ T & \phi / \nu & \Xi - \phi \theta / \nu \end{array}$$

and satisfying  $(\epsilon_x; \epsilon_u; \rho_{ux}^\pm) \xrightarrow{\beta} \left( \frac{1}{u} \middle| x; \frac{1}{u} \middle| ; \frac{1}{u} \middle| T_u^{\pm 1} - 1 \right)$ .


**Proposition.** If  $T$  is a u-tangle and  $\beta(\delta T) = (\omega, A)$ , then  $\gamma(T) = (\omega, \sigma - A)$ , where  $\sigma = \text{diag}(\sigma_a)_{a \in S}$ . Under this,  $m_c^{ab} \leftrightarrow tha^{ab} // tm_c^{ab} // hm_c^{ab}$ .

**References.**


[BN] D. Bar-Natan, *Balloons and Hoops and their Universal Finite Type Invariant, BF Theory, and an Ultimate Alexander Invariant*, ωεβ/KBH, arXiv:1308.1721.  
 [BND] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects I-II*, ωεβ/WKO1, ωεβ/WKO2, arXiv:1405.1956, arXiv:1405.1955.  
 [BNS] D. Bar-Natan and S. Selmani, *Meta-Monoids, Meta-Bicrossed Products, and the Alexander Polynomial*, J. of Knot Theory and its Ramifications **22-10** (2013), arXiv:1302.5689.  
 [CR] A. S. Cattaneo and C. A. Rossi, *Wilson Surfaces and Higher Dimensional Knot Invariants*, Commun. in Math. Phys. **256-3** (2005) 513–537, arXiv:math-ph/0210037.  
 [CT] D. Cimasoni and V. Turaev, *A Lagrangian Representation of Tangles*, Topology **44** (2005) 747–767, arXiv:math.GT/0406269.  
 [KLW] P. Kirk, C. Livingston, and Z. Wang, *The Gassner Representation for String Links*, Comm. Cont. Math. **3** (2001) 87–136, arXiv:math/9806035.  
 [LD] J. Y. Le Dimet, *Enlacements d'Intervalles et Représentation de Gassner*, Comment. Math. Helv. **67** (1992) 306–315.

**Theorem 3 [BND, BN].**  $\exists!$  a homomorphic expansion, aka a homomorphic universal finite type invariant  $Z$  of  $w$ -knotted balloons and hoops.  $\zeta := \log Z$  takes values in  $FL(T)^H \times CW(T)$ .

$\zeta$  is computable!  $\zeta$  of the Borromean tangle, to degree 5:



**Proposition [BN].** Modulo all relations that universally hold for the 2D non-Abelian Lie algebra and after some changes-of-variable,  $\zeta$  reduces to  $\beta$  and the KBH operations on  $\zeta$  reduce to the formulas in Theorem 2.



**A Big Question.** Does it all extend to arbitrary 2-knots (not necessarily “simple”)? To arbitrary codimension-2 knots?

**BF Following [CR].**  $A \in \Omega^1(M = \mathbb{R}^4, \mathfrak{g})$ ,  $B \in \Omega^2(M, \mathfrak{g}^*)$ ,

$$S(A, B) := \int_M \langle B, F_A \rangle.$$


With  $\kappa: (S = \mathbb{R}^2) \rightarrow M$ ,  $\beta \in \Omega^0(S, \mathfrak{g})$ ,  $\alpha \in \Omega^1(S, \mathfrak{g}^*)$ , set

$$O(A, B, \kappa) := \int \mathcal{D}\beta \mathcal{D}\alpha \exp\left(\frac{i}{\hbar} \int_S \langle \beta, d_\kappa \alpha + \kappa^* B \rangle\right).$$

**The BF Feynman Rules.** For an edge  $e$ , let  $\Phi_e$  be its direction, in  $S^3$  or  $S^1$ . Let  $\omega_3$  and  $\omega_1$  be volume forms on  $S^3$  and  $S^1$ . Then

$$Z_{BF} = \sum_{\text{diagrams } D} \frac{|D|}{|\text{Aut}(D)|} \int_{\mathbb{R}^2} \dots \int_{\mathbb{R}^2} \int_{\mathbb{R}^4} \dots \int_{\mathbb{R}^4} \prod_{\text{red } e \in D} \Phi_e^* \omega_3 \prod_{\text{black } e \in D} \Phi_e^* \omega_1$$

(modulo some  $STU$ - and  $IHX$ -like relations).



**Issues.** • Signs don't quite work out, and BF seems to reproduce only “half” of the wheels invariant.

• There are many more configuration space integrals than BF Feynman diagrams and than just trees and wheels.

• I don't know how to define “finite type” for arbitrary 2-knots.



“God created the knots, all else in topology is the work of mortals.”

Leopold Kronecker (modified)



# Meta-Groups, Meta-Bicrossed-Products, and the Alexander Polynomial, 1



Dror Bar-Natan in Montreal, June 2013.

<http://www.math.toronto.edu/~drorbn/Talks/Montreal-1306/>

**Abstract.** I will define “meta-groups” and explain how one specific meta-group, which in itself is a “meta-bicrossed-product”, gives rise to an “ultimate Alexander invariant” of tangles, that contains the Alexander polynomial (multivariable, if you wish), has extremely good composition properties, is evaluated in a topologically meaningful way, and is least-wasteful in a computational sense. If you believe in categorification, that’s a wonderful playground.



**This work is closely related** to work by Le Dimet (Comment. Math. Helv. **67** (1992) 306–315), Kirk, Livingston and Wang (arXiv:math/9806035) and Cimasoni and Turaev (arXiv:math.GT/0406269).

**See also** Dror Bar-Natan and Sam Selmani, *Meta-Monoids, Meta-Bicrossed Products, and the Alexander Polynomial*, arXiv:1302.5689.

## Alexander Issues.


- Quick to compute, but computation departs from topology.
- Extends to tangles, but at an exponential cost.
- Hard to categorify.

**Idea.** Given a group  $G$  and two “YB” pairs  $R^\pm = (g_o^\pm, g_u^\pm) \in G^2$ , map them to xings and “multiply along”, so that



**This Fails!** R2 implies that  $g_o^\pm g_o^\mp = e = g_u^\pm g_u^\mp$  and then R3 implies that  $g_o^\pm$  and  $g_u^\pm$  commute, so the result is a simple counting invariant.



**A Group Computer.** Given  $G$ , can store group elements and perform operations on them:



Also has  $S_x$  for inversion,  $e_x$  for unit insertion,  $d_x$  for register deletion,  $\Delta_{xy}^z$  for element cloning,  $\rho_y^x$  for renamings, and  $(D_1, D_2) \mapsto D_1 \cup D_2$  for merging, and many obvious composition axioms relating those.

$$P = \{x : g_1, y : g_2\} \Rightarrow P = \{d_y P\} \cup \{d_x P\}$$

**Riddle.** People often study  $\pi_1(X) = [S^1, X]$  and  $\pi_2(X) = [S^2, X]$ . Why not  $\pi_T(X) := [T, X]$ ?



**A Meta-Group.** Is a similar “computer”, only its internal structure is unknown to us. Namely it is a collection of sets  $\{G_\gamma\}$  indexed by all finite sets  $\gamma$ , and a collection of operations  $m_z^{xy}$ ,  $S_x$ ,  $e_x$ ,  $d_x$ ,  $\Delta_{xy}^z$  (sometimes),  $\rho_y^x$ , and  $\cup$ , satisfying the exact same *linear* properties.

**Example 0.** The non-meta example,  $G_\gamma := G^\gamma$ .


**Example 1.**  $G_\gamma := M_{\gamma \times \gamma}(\mathbb{Z})$ , with simultaneous row and column operations, and “block diagonal” merges. Here if

$P = \begin{pmatrix} x & a & b \\ y & c & d \end{pmatrix}$  then  $d_y P = (x : a)$  and  $d_x P = (y : d)$  so

$\{d_y P\} \cup \{d_x P\} = \begin{pmatrix} x & a & 0 \\ y & 0 & d \end{pmatrix} \neq P$ . So this  $G$  is truly meta.

**Claim.** From a meta-group  $G$  and YB elements  $R^\pm \in G_2$  we can construct a knot/tangle invariant.


**A Standard Alexander Formula.** Label the arcs  $1$  through  $(n + 1) = 1$ , make an  $n \times n$  matrix as below, delete one row and one column, and compute the determinant:



$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & x-1 & 0 & -x \\ -1 & x & 0 & 0 & 0 & 0 & 0 & 1-x & 0 \\ 0 & -1 & x & 0 & 1-x & 0 & 0 & 0 & 0 \\ x-1 & 0 & -x & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1-x & 0 & -1 & x & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -x & 1 & 0 & 0 & x-1 \\ 0 & 0 & 1-x & 0 & 0 & -1 & x & 0 & 0 \\ 0 & 0 & 0 & x-1 & 0 & 0 & -x & 1 & 0 \end{pmatrix} \quad [[1 ;; 7, 1 ;; 7]] // \text{Det}$$

$$-1 + 4x - 8x^2 + 11x^3 - 8x^4 + 4x^5 - x^6$$

**Bicrossed Products.** If  $G = HT$  is a group presented as a product of two of its subgroups, with  $H \cap T = \{e\}$ , then also  $G = TH$  and  $G$  is determined by  $H$ ,  $T$ , and the “swap” map  $sw^{th} : (t, h) \mapsto (h', t')$  defined by  $th = h't'$ . The map  $sw$  satisfies (1) and (2) below; conversely, if  $sw : T \times H \rightarrow H \times T$  satisfies (1) and (2) (+ lesser conditions), then (3) defines a group structure on  $H \times T$ , the “bicrossed product”.



# Meta-Groups, Meta-Bicrossed-Products, and the Alexander Polynomial, 2

A **Meta-Bicrossed-Product** is a collection of sets  $\beta(\eta, \tau)$  and operations  $tm_w^{uv}$ ,  $hm_z^{xy}$  and  $sw_{ux}^{th}$  (and lesser ones), such that  $tm$  and  $hm$  are “associative” and (1) and (2) hold (+ lesser conditions). A meta-bicrossed-product defines a meta-group with  $G_\gamma := \beta(\gamma, \gamma)$  and  $gm$  as in (3).

**Example.** Take  $\beta(\eta, \tau) = M_{\tau \times \eta}(\mathbb{Z})$  with row operations for the tails, column operations for the heads, and a trivial swap.

**$\beta$  Calculus.** Let  $\beta(\eta, \tau)$  be

$$\left\{ \begin{array}{c|ccc} \omega & h_1 & h_2 & \cdots \\ t_1 & \alpha_{11} & \alpha_{12} & \cdot \\ t_2 & \alpha_{21} & \alpha_{22} & \cdot \\ \vdots & \cdot & \cdot & \cdot \end{array} \middle| \begin{array}{l} h_j \in \eta, t_i \in \tau, \text{ and } \omega \text{ and} \\ \text{the } \alpha_{ij} \text{ are rational func-} \\ \text{tions in a variable } X \text{ with} \\ \omega(1) = 1 \text{ and } \alpha_{ij}(1) = 0 \end{array} \right\},$$

$$tm_w^{uv} : \begin{array}{c|c} \omega & \cdots \\ t_u & \alpha \\ t_v & \beta \\ \vdots & \gamma \end{array} \mapsto \begin{array}{c|c} \omega & \cdots \\ t_w & \alpha + \beta \\ \vdots & \gamma \end{array}, \quad \begin{array}{c|c} \omega_1 & \eta_1 \\ \tau_1 & \alpha_1 \end{array} \cup \begin{array}{c|c} \omega_2 & \eta_2 \\ \tau_2 & \alpha_2 \end{array} = \begin{array}{c|c} \omega_1\omega_2 & \eta_1 \eta_2 \\ \tau_1 & \alpha_1 \ 0 \\ \tau_2 & 0 \ \alpha_2 \end{array},$$

$$hm_z^{xy} : \begin{array}{c|ccc} \omega & h_x & h_y & \cdots \\ \vdots & \alpha & \beta & \gamma \end{array} \mapsto \begin{array}{c|ccc} \omega & h_z & & \cdots \\ \vdots & \alpha + \beta + \langle \alpha \rangle \beta & & \gamma \end{array},$$

$$sw_{ux}^{th} : \begin{array}{c|ccc} \omega & h_x & \cdots \\ t_u & \alpha & \beta \\ \vdots & \gamma & \delta \end{array} \mapsto \begin{array}{c|ccc} \omega\epsilon & h_x & \cdots \\ t_u & \alpha(1 + \langle \gamma \rangle / \epsilon) & \beta(1 + \langle \gamma \rangle / \epsilon) \\ \vdots & \gamma / \epsilon & \delta - \gamma\beta / \epsilon \end{array},$$

where  $\epsilon := 1 + \alpha$  and  $\langle c \rangle := \sum_i c_i$ , and let

$$R_{ab}^p := \begin{array}{c|cc} 1 & h_a & h_b \\ t_a & 0 & X - 1 \\ t_b & 0 & 0 \end{array} \quad R_{ab}^m := \begin{array}{c|cc} 1 & h_a & h_b \\ t_a & 0 & X^{-1} - 1 \\ t_b & 0 & 0 \end{array}.$$

**Theorem.**  $Z^\beta$  is a tangle invariant (and more). Restricted to knots, the  $\omega$  part is the Alexander polynomial. On braids, it is equivalent to the Burau representation. A variant for links contains the multivariable Alexander polynomial.

**Why Happy?** • Applications to w-knots.

• Everything that I know about the Alexander polynomial can be expressed cleanly in this language (even if without proof), except HF, but including genus, ribboness, cabling, v-knots, knotted graphs, etc., and there’s potential for vast generalizations.

• The least wasteful “Alexander for tangles” I’m aware of.

• Every step along the computation is the invariant of something.

• Fits on one sheet, including implementation & propaganda.



**Further meta-monoids.**  $\Pi$  (and variants),  $\mathcal{A}$  (and quotients),  $vT$ , ...

**Further meta-bicrossed-products.**  $\Pi$  (and variants),  $\vec{\mathcal{A}}$  (and quotients),  $M_0$ ,  $M$ ,  $\mathcal{K}^{bh}$ ,  $\mathcal{K}^{rbh}$ , ...

**Meta-Lie-algebras.**  $\mathcal{A}$  (and quotients),  $\mathcal{S}$ , ...

**Meta-Lie-bialgebras.**  $\vec{\mathcal{A}}$  (and quotients), ...

I don’t understand the relationship between  $gr$  and  $H$ , as it appears, for example, in braid theory.

I mean business!

```


SSimp = Factor; SetAttributes[BCollect, Listable];
BCollect[B[omega_, A_] := B[SSimp[omega],
Collect[A, h, Collect[omega, t, SSimp] &];
BForm[B[omega_, A_] := Module[{ts, hs, M},
ts = Union[Cases[B[omega, A], t_u -> u, Infinity]];
hs = Union[Cases[B[omega, A], h_x -> x, Infinity]];
M = Outer[SSimp[Coefficient[A, h_x t_u]] &, hs, ts];
PrependTo[M, t_u & /@ ts];
M = Prepend[Transpose[M], Prepend[h_x & /@ hs, omega]];
MatrixForm[M];
BForm[else_] := else /. B_B -> BForm[B];
Format[B_B, StandardForm] := BForm[B];
    
```

```

<omega> := # /. t_u -> 1;
tm_u_v_w[omega_] := BCollect[B /. t_u | v -> t_u];
hm_x_y_z[B[omega_, A_] := Module[
{alpha = D[A, h_x], beta = D[A, h_y], gamma = A /. h_x | y -> 0},
B[omega, (alpha + (1 + alpha) beta) h_x + gamma] // BCollect;
sw_u_x[B[omega_, A_] := Module[{alpha, beta, gamma, delta, epsilon},
alpha = Coefficient[A, h_x t_u]; beta = D[A, t_u] /. h_x -> 0;
gamma = D[A, h_x] /. t_u -> 0; delta = A /. h_x | t_u -> 0;
epsilon = 1 + alpha;
B[omega + epsilon, alpha (1 + gamma / epsilon) h_x t_u + beta (1 + gamma / epsilon) t_u
+ gamma / epsilon h_x + delta - gamma * beta / epsilon
] // BCollect;
];
gm_u_v_w[omega_] := beta // sw_ab // hm_u_v_w // tm_u_v_w;
B /. B[omega_, A_] B[omega_2_, A2_] := B[omega_1 * omega_2, A1 + A2];
Rp_a_b := B[1, (X - 1) t_u h_x];
Rm_a_b := B[1, (X^-1 - 1) t_u h_x];
    
```

$$\{\beta = \mathbf{B}[\omega, \text{Sum}[\alpha_{10+i+j} t_i h_j, \{\mathbf{i}, \{1, 2, 3\}\}, \{\mathbf{j}, \{4, 5\}\}]\},$$

$$(\beta // tm_{12 \rightarrow 1} // sw_{14}) = (\beta // sw_{24} // sw_{14} // tm_{12 \rightarrow 1})$$



$$\left\{ \begin{array}{c} \omega \quad h_4 \quad h_5 \\ t_1 \quad \alpha_{14} \quad \alpha_{15} \\ t_2 \quad \alpha_{24} \quad \alpha_{25} \\ t_3 \quad \alpha_{34} \quad \alpha_{35} \end{array} \right\}, \text{True}$$

$$\left\{ \begin{array}{ccc} 1 & h_1 & h_2 \\ t_2 & -\frac{-1+X}{X} & 0 \\ t_3 & -\frac{-1+X}{X} & -\frac{-1+X}{X} \end{array} \right\}, \left\{ \begin{array}{ccc} 1 & h_1 & h_2 \\ t_2 & -\frac{-1+X}{X} & 0 \\ t_3 & -\frac{-1+X}{X} & -\frac{-1+X}{X} \end{array} \right\}$$

... divide and conquer!

$$\beta = Rm_{12,1} Rm_{27} Rm_{83} Rm_{4,11} Rp_{16,5} Rp_{6,13} Rp_{14,9} Rp_{10,15}$$


$$\begin{pmatrix} 1 & h_1 & h_3 & h_5 & h_7 & h_9 & h_{11} & h_{13} & h_{15} \\ t_2 & 0 & 0 & 0 & -\frac{-1+X}{X} & 0 & 0 & 0 & 0 \\ t_4 & 0 & 0 & 0 & 0 & 0 & -\frac{-1+X}{X} & 0 & 0 \\ t_6 & 0 & 0 & 0 & 0 & 0 & 0 & -1+X & 0 \\ t_8 & 0 & -\frac{-1+X}{X} & 0 & 0 & 0 & 0 & 0 & 0 \\ t_{10} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1+X \\ t_{12} & -\frac{-1+X}{X} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ t_{14} & 0 & 0 & 0 & 0 & -1+X & 0 & 0 & 0 \\ t_{16} & 0 & 0 & -1+X & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Do[ $\beta = \beta // gm_{1k \rightarrow 1}, \{k, 2, 10\}$ ];  $\beta$




$$\begin{pmatrix} \frac{1}{X} & h_1 & & h_{11} & & h_{13} & & h_{15} \\ t_1 & -\frac{(-1+X)(1+X)}{X} & & (-1+X)(1-X+X^2) & & (-1+X)(1-X+X^2) & & -1+X \\ t_{12} & -\frac{-1+X}{X} & & 0 & & 0 & & 0 \\ t_{14} & -1+X & & \frac{(-1+X)^2(1-X+X^2)}{X} & & -\frac{(-1+X)^2(1-X+X^2)}{X} & & 0 \\ t_{16} & -\frac{-1+X}{X} & & (-1+X)^2 & & -\frac{(-1+X)^3}{X} & & 0 \end{pmatrix}$$

Do[ $\beta = \beta // gm_{1k \rightarrow 1}, \{k, 11, 16\}$ ];  $\beta$

$$\left( -\frac{1-4X+8X^2-11X^3+8X^4-4X^5+X^6}{X^3} \right)$$



1. Where does it *more simply* come from?
2. Remove all the denominators.
3. How do determinants arise in this context?
4. Understand links (“meta-conjugacy classes”).
5. Find the “reality condition”.
6. Do some “Algebraic Knot Theory”.
7. Categorify.
8. Do the same in other natural quotients of the v/w-story.



"God created the knots, all else in topology is the work of mortals."  
 Leopold Kronecker (modified)

www.katlas.org The Knot Atlas  
 Inverse Car Talk

example

# Trees and Wheels and Balloons and Hoops

Dror Bar-Natan, Zurich, September 2013

$\omega\epsilon\beta$ : <http://www.math.toronto.edu/~drorbn/Talks/Zurich-130919>



## 15 Minutes on Algebra

Let  $T$  be a finite set of “tail labels” and  $H$  a finite set of “head labels”. Set

$$M_{1/2}(T; H) := FL(T)^H,$$

“ $H$ -labeled lists of elements of the degree-completed free Lie algebra generated by  $T$ ”.

$$FL(T) = \left\{ 2t_2 - \frac{1}{2}[t_1, [t_1, t_2]] + \dots \right\} / \left( \begin{array}{c} \text{anti-symmetry} \\ \text{Jacobi} \end{array} \right)$$

... with the obvious bracket.

$$M_{1/2}(u, v; x, y) = \left\{ \lambda = \left( x \rightarrow \begin{array}{c} u \quad v \\ \diagdown \quad \diagup \\ x \end{array}, y \rightarrow \begin{array}{c} v \\ \downarrow \\ y \end{array} - \frac{22}{7} \begin{array}{c} u \quad v \\ \diagdown \quad \diagup \\ y \end{array} \right) \dots \right\}$$

Operations  $M_{1/2} \rightarrow M_{1/2}$ . ↙ newspeak!

**Tail Multiply**  $tm_w^{uv}$  is  $\lambda \mapsto \lambda \parallel (u, v \rightarrow w)$ , satisfies “meta-associativity”,  $tm_u^{uv} \parallel tm_u^{vw} = tm_v^{uv} \parallel tm_v^{vw}$ .

**Head Multiply**  $hm_z^{xy}$  is  $\lambda \mapsto (\lambda \setminus \{x, y\}) \cup (z \rightarrow \text{bch}(\lambda_x, \lambda_y))$ , where

$$\text{bch}(\alpha, \beta) := \log(e^\alpha e^\beta) = \alpha + \beta + \frac{[\alpha, \beta]}{2} + \frac{[\alpha, [\alpha, \beta]] + [[\alpha, \beta], \beta]}{12} + \dots$$

satisfies  $\text{bch}(\text{bch}(\alpha, \beta), \gamma) = \log(e^{\alpha} e^{\beta} e^{\gamma}) = \text{bch}(\alpha, \text{bch}(\beta, \gamma))$  and hence meta-associativity,  $hm_x^{xy} \parallel hm_x^{yz} = hm_y^{xy} \parallel hm_y^{yz}$ .

**Tail by Head Action**  $tha^{ux}$  is  $\lambda \mapsto \lambda \parallel RC_u^{\lambda_x}$ , where  $C_u^{-\gamma}: FL \rightarrow FL$  is the substitution  $u \rightarrow e^{-\gamma} u e^{\gamma}$ , or more precisely,





$$C_u^{-\gamma}: u \rightarrow e^{-\text{ad} \gamma}(u) = u - [\gamma, u] + \frac{1}{2}[\gamma, [\gamma, u]] - \dots,$$

and  $RC_u^{\gamma} = (C_u^{-\gamma})^{-1}$ . Then  $C_u^{\text{bch}(\alpha, \beta)} = C_u^{\alpha} \parallel RC_u^{-\beta} \parallel C_u^{\beta}$  hence  $RC_u^{\text{bch}(\alpha, \beta)} = RC_u^{\alpha} \parallel RC_u^{\beta} \parallel RC_u^{\alpha}$  hence “meta  $u^{xy} = (u^x)^y$ ”,

$$hm_z^{xy} \parallel tha^{uz} = tha^{ux} \parallel tha^{uy} \parallel hm_z^{xy},$$

and  $tm_w^{uv} \parallel C_w^{\gamma} \parallel tm_w^{uv} = C_u^{\gamma} \parallel RC_u^{-\gamma} \parallel C_u^{\gamma} \parallel tm_w^{uv}$  and hence “meta  $(uv)^x = u^x v^x$ ”,  $tm_w^{uv} \parallel tha^{wx} = tha^{ux} \parallel tha^{vx} \parallel tm_w^{uv}$ .


**Wheels.** Let  $M(T; H) := M_{1/2}(T; H) \times CW(T)$ , where  $CW(T)$  is the (completed graded) vector space of cyclic words on  $T$ , or equally well, on  $FL(T)$ :





**Operations.** On  $M(T; H)$ , define  $tm_w^{uv}$  and  $hm_z^{xy}$  as before, and  $tha^{ux}$  by adding some  $J$ -spice:

$$(\lambda; \omega) \mapsto (\lambda, \omega + J_u(\lambda_x)) \parallel RC_u^{\lambda_x},$$

where  $J_u(\gamma) := \int_0^1 ds \text{div}_u(\gamma \parallel RC_u^{s\gamma}) \parallel C_u^{-s\gamma}$ , and



$\text{div}_u \rightarrow$




**Theorem Blue.** All blue identities still hold.

**Merge Operation.**  $(\lambda_1; \omega_1) * (\lambda_2; \omega_2) := (\lambda_1 \cup \lambda_2; \omega_1 + \omega_2)$ .


$\mathcal{K}^{bh}(T; H)$ .

## 15 Minutes on Topology

“Ribbon-knotted balloons and hoops”




**Examples.**



**More on  $\delta$**



satisfies R123, VR123, D, and




- $\delta$  injects u-knots into  $\mathcal{K}^{bh}$  (likely u-tangles too).
- $\delta$  maps v-tangles to  $\mathcal{K}^{bh}$ ; the kernel contains the above and **conjecturally** (Sato), that's all.
- Allowing punctures and cuts,  $\delta$  is onto.

**Operations**


Punctures & Cuts

Connected Sums.




If  $X$  is a space,  $\pi_1(X)$  is a group,  $\pi_2(X)$  is an Abelian group, and  $\pi_1$  acts on  $\pi_2$ .




**Riddle.** People often study  $\pi_1(X) = [S^1, X]$  and  $\pi_2(X) = [S^2, X]$ . Why not  $\pi_T(X) := [T, X]$ ?



$K$ :



$K \parallel hm_z^{xy}$ :




“Meta-Group-Action”


**Properties.**

- Associativities:  $m_a^{ab} \parallel m_a^{ac} = m_b^{bc} \parallel m_a^{ab}$ , for  $m = tm, hm$ .
- “( $uv$ )<sup>x</sup> =  $u^x v^x$ ”:  $tm_w^{uv} \parallel tha^{wx} = tha^{ux} \parallel tha^{vx} \parallel tm_w^{uv}$ ,
- “( $u(xy)$ ) = ( $u^x$ )<sup>y</sup>”:  $hm_z^{xy} \parallel tha^{uz} = tha^{ux} \parallel tha^{uy} \parallel hm_z^{xy}$ .

**Tangle concatenations**  $\rightarrow \pi_1 \times \pi_2$ . With  $dm_c^{ab} := tha^{ab} \parallel tm_c^{ab} \parallel hm_c^{ab}$ ,



**Finite type** invariants make sense in the usual way, and



“algebra” is (the primitive part of) “gr” of “topology”.

# Trees and Wheels and Balloons and Hoops: Why I Care

**Moral.** To construct an  $M$ -valued invariant  $\zeta$  of  $(v-)$ tangles, and nearly an invariant on  $\mathcal{K}^{bh}$ , it is enough to declare  $\zeta$  on the generators, and verify the relations that  $\delta$  satisfies.

**The Invariant  $\zeta$ .** Set  $\zeta(\epsilon_x) = (x \rightarrow 0; 0)$ ,  $\zeta(\epsilon_u) = (( ); 0)$ , and

$$\zeta: \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} \mapsto \begin{array}{c} \left( \begin{array}{c} u \\ \downarrow x \end{array}; 0 \right) \\ \left( - \begin{array}{c} u \\ \downarrow x \end{array}; 0 \right) \end{array}$$

**Theorem.**  $\zeta$  is (log of) the unique homomorphic universal finite type invariant on  $\mathcal{K}^{bh}$ .  
 (... and is the tip of an iceberg)

Paper in progress with Danco,  $\omega\epsilon\beta/wko$



See also  $\omega\epsilon\beta/tenn$ ,  $\omega\epsilon\beta/bonn$ ,  $\omega\epsilon\beta/swiss$ ,  $\omega\epsilon\beta/portfolio$

**$\zeta$  is computable!**  $\zeta$  of the Borromean tangle, to degree 5:

**Tensorial Interpretation.** Let  $\mathfrak{g}$  be a finite dimensional Lie algebra (any!). Then there's  $\tau : FL(T) \rightarrow \text{Fun}(\oplus_T \mathfrak{g} \rightarrow \mathfrak{g})$  and  $\tau : CW(T) \rightarrow \text{Fun}(\oplus_T \mathfrak{g})$ . Together,  $\tau : M(T; H) \rightarrow \text{Fun}(\oplus_T \mathfrak{g} \rightarrow \oplus_H \mathfrak{g})$ , and hence

$$e^\tau : M(T; H) \rightarrow \text{Fun}(\oplus_T \mathfrak{g} \rightarrow \mathcal{U}^{\otimes H}(\mathfrak{g})).$$

**$\zeta$  and BF Theory.** (See Cattaneo-Rossi, arXiv:math-ph/0210037) Let  $A$  denote a  $\mathfrak{g}$ -connection on  $S^4$  with curvature  $F_A$ , and  $B$  a  $\mathfrak{g}^*$ -valued 2-form on  $S^4$ . For a hoop  $\gamma_x$ , let  $\text{hol}_{\gamma_x}(A) \in \mathcal{U}(\mathfrak{g})$  be the holonomy of  $A$  along  $\gamma_x$ . For a ball  $\gamma_u$ , let  $\mathcal{O}_{\gamma_u}(B) \in \mathfrak{g}^*$  be (roughly) the integral of  $B$  (transported via  $A$  to  $\infty$ ) on  $\gamma_u$ .




**Loose Conjecture.** For  $\gamma \in \mathcal{K}(T; H)$ ,

$$\int \mathcal{D}A \mathcal{D}B e^{\int B \wedge F_A} \prod_u e^{\mathcal{O}_{\gamma_u}(B)} \bigotimes_x \text{hol}_{\gamma_x}(A) = e^\tau(\zeta(\gamma)).$$

That is,  $\zeta$  is a complete evaluation of the BF TQFT.

"God created the knots, all else in topology is the work of mortals."  
 Leopold Kronecker (modified) [www.katlas.org](http://www.katlas.org) The Knot Atlas

The  $\beta$  quotient is  $M$  divided by all relations that universally hold when  $\mathfrak{g}$  is the 2D non-Abelian Lie algebra. Let  $R = \mathbb{Q}[\{c_u\}_{u \in T}]$  and  $L_\beta := R \otimes T$  with central  $R$  and with  $[u, v] = c_u v - c_v u$  for  $u, v \in T$ . Then  $FL \rightarrow L_\beta$  and  $CW \rightarrow R$ . Under this,



$$\mu \rightarrow ((\lambda_x); \omega) \quad \text{with } \lambda_x = \sum_{u \in T} \lambda_{ux} u x, \quad \lambda_{ux}, \omega \in R,$$

$$\text{bch}(u, v) \rightarrow \frac{c_u + c_v}{e^{c_u + c_v} - 1} \left( \frac{e^{c_u} - 1}{c_u} u + e^{c_u} \frac{e^{c_v} - 1}{c_v} v \right),$$


if  $\gamma = \sum \gamma_v v$  then with  $c_\gamma := \sum \gamma_v c_v$ ,

$$u \parallel RC_\gamma^u = \left( 1 + c_u \gamma_u \frac{e^{c_\gamma} - 1}{c_\gamma} \right)^{-1} \left( e^{c_\gamma} u - c_u \frac{e^{c_\gamma} - 1}{c_\gamma} \sum_{v \neq u} \gamma_v v \right),$$

$\text{div}_u \gamma = c_u \gamma_u$ , and  $J_u(\gamma) = \log \left( 1 + \frac{e^{c_\gamma} - 1}{c_\gamma} c_u \gamma_u \right)$ , so  $\zeta$  is formula-computable to all orders! **Can we simplify?**

**Repackaging.** Given  $((x \rightarrow \lambda_{ux}); \omega)$ , set  $c_x := \sum_v c_v \lambda_{vx}$ , replace  $\lambda_{ux} \rightarrow \alpha_{ux} := c_u \lambda_{ux} \frac{e^{c_x} - 1}{c_x}$  and  $\omega \rightarrow e^\omega$ , use  $t_u = e^{c_u}$ , and write  $\alpha_{ux}$  as a matrix. Get " **$\beta$  calculus**".

**$\beta$  Calculus.** Let  $\beta(T; H)$  be

$$\left\{ \begin{array}{c|ccc} \omega & x & y & \cdots \\ u & \alpha_{ux} & \alpha_{uy} & \cdot \\ v & \alpha_{vx} & \alpha_{vy} & \cdot \\ \vdots & \cdot & \cdot & \cdot \end{array} \middle| \begin{array}{l} \omega \text{ and the } \alpha_{ux} \text{'s are} \\ \text{rational functions in} \\ \text{variables } t_u, \text{ one for} \\ \text{each } u \in T. \end{array} \right\},$$


$$tm_w^{uv} : \begin{array}{c|cc} \omega & \cdots \\ u & \alpha \\ v & \beta \\ \vdots & \gamma \end{array} \mapsto \begin{array}{c|cc} \omega & \cdots \\ w & \alpha + \beta \\ & \gamma \end{array}, \quad \begin{array}{c|cc} \omega_1 & H_1 & \omega_2 & H_2 \\ T_1 & \alpha_1 & T_2 & \alpha_2 \\ \hline & \omega_1 \omega_2 & H_1 & H_2 \\ & T_1 & \alpha_1 & 0 \\ & T_2 & 0 & \alpha_2 \end{array},$$

$$hm_z^{xy} : \begin{array}{c|ccc} \omega & x & y & \cdots \\ \vdots & \alpha & \beta & \gamma \end{array} \mapsto \begin{array}{c|cc} \omega & z & \cdots \\ \vdots & \alpha + \beta + \langle \alpha \rangle \beta & \gamma \end{array},$$

$$tha_{ux} : \begin{array}{c|ccc} \omega & x & \cdots \\ u & \alpha & \beta \\ \vdots & \gamma & \delta \end{array} \mapsto \begin{array}{c|cc} \omega \epsilon & x & \cdots \\ u & \alpha(1 + \langle \gamma \rangle / \epsilon) & \beta(1 + \langle \gamma \rangle / \epsilon) \\ \vdots & \gamma / \epsilon & \delta - \gamma \beta / \epsilon \end{array},$$

where  $\epsilon := 1 + \alpha$ ,  $\langle \alpha \rangle := \sum_v \alpha_v$ , and  $\langle \gamma \rangle := \sum_{v \neq u} \gamma_v$ , and let

$$R_{ux}^+ := \frac{1}{u} \left| \begin{array}{c} x \\ t_u - 1 \end{array} \right. \quad R_{ux}^- := \frac{1}{u} \left| \begin{array}{c} x \\ t_u^{-1} - 1 \end{array} \right.$$

On long knots,  $\omega$  is the Alexander polynomial!

**Why happy?** An ultimate Alexander invariant: Manifestly polynomial (time and size) extension of the (multivariable) Alexander polynomial to tangles. Every step of the computation is the computation of the invariant of some topological thing (no fishy Gaussian elimination). *If there should be an Alexander invariant with a computable algebraic categorification, it is this one!*




See also  $\omega\epsilon\beta/regina$ ,  $\omega\epsilon\beta/caen$ ,  $\omega\epsilon\beta/newton$ .

**May class:**  $\omega\epsilon\beta/aarhus$  **Class next year:**  $\omega\epsilon\beta/1350$   
**Paper:**  $\omega\epsilon\beta/kbh$





# Finite Type Invariants of Ribbon Knotted Balloons and Hoops

**Abstract.** On my September 17 Geneva talk ( $\omega/\text{sep}$ ) I described a certain trees-and-wheels-valued invariant  $\zeta$  of ribbon knotted loops and 2-spheres in 4-space, and my October 8 Geneva talk ( $\omega/\text{oct}$ ) describes its reduction to the Alexander polynomial. Today I will explain how that same invariant arises completely naturally within the theory of finite type invariants of ribbon knotted loops and 2-spheres in 4-space.



My goal is to tell you why such an invariant is expected, yet not to derive the computable formulas.



**Expansions**  
the semi-virtual  $\otimes := \diagup - \diagdown$  i.e.  $\diagup - \diagdown$  or  $\diagdown - \diagup$

Let  $\mathcal{I}^n := \langle \text{pictures with } \geq n \text{ semi-virts} \rangle \subset \mathcal{K}^{bh}$ .  
We seek an "expansion"

$$Z: \mathcal{K}^{bh} \rightarrow \text{gr } \mathcal{K}^{bh} = \widehat{\bigoplus} \mathcal{I}^n / \mathcal{I}^{n+1} =: \mathcal{A}^{bh}$$

satisfying "property U": if  $\gamma \in \mathcal{I}^n$ , then


$$Z(\gamma) = (0, \dots, 0, \gamma / \mathcal{I}^{n+1}, *, *, \dots).$$



X.-S. Lin

**Why?** • Just because, and this is vastly more general.  
•  $(\mathcal{K}^{bh} / \mathcal{I}^{n+1})^*$  is "finite-type/polynomial invariants".  
• The Taylor example: Take  $\mathcal{K} = C^\infty(\mathbb{R}^n)$ ,  $\mathcal{I} = \{f \in \mathcal{K} : f(0) = 0\}$ . Then  $\mathcal{I}^n = \{f : f \text{ vanishes like } |x|^n\}$  so  $\mathcal{I}^n / \mathcal{I}^{n+1}$  is homogeneous polynomials of degree  $n$  and  $Z$  is a "Taylor expansion"! (So Taylor expansions are vastly more general than you'd think).

**Plan.** We'll construct a graded  $\tilde{\mathcal{A}}^{bh}$ , a surjective graded  $\pi: \tilde{\mathcal{A}}^{bh} \rightarrow \mathcal{A}^{bh}$ , and a filtered  $\tilde{Z}: \mathcal{K}^{bh} \rightarrow \tilde{\mathcal{A}}^{bh}$  so that  $\pi \circ \text{gr } \tilde{Z} = \text{Id}$  (property U: if  $\text{deg } D = n$ ,  $\tilde{Z}(\pi(D)) = \pi(D) + (\text{deg } \geq n)$ ). Hence •  $\pi$  is an isomorphism. •  $Z := \tilde{Z} \circ \pi$  is an expansion.




"God created the knots, all else in topology is the work of mortals."

Leopold Kronecker (modified)


www.katlas.org




## Action 1.




## Deriving 4T.



## Action 2.



## The Bracket-Rise Theorem.




## Proof.



**Corollaries.** (1) Related to Lie algebras! (2) Only trees and wheels persist.

**Theorem.**  $\mathcal{A}^{bh}$  is a bi-algebra. The space of its primitives is  $FL(T)^H \times CW(T)$ , and  $\zeta = \log Z$ .

$\zeta$  is computable!  $\zeta$  of the Borromean tangle, to degree 5:



I have a nice free-Lie calculator!



# The Kashiwara-Vergne Problem and Topology

**Abstract.** I will describe a general machine, a close cousin of Taylor's theorem, whose inputs are topics in topology and whose outputs are problems in algebra. There are many inputs the machine can take, and many outputs it produces, but I will concentrate on just one input/output pair. When fed with a certain class of knotted 2-dimensional objects in 4-dimensional space, it outputs the Kashiwara-Vergne Problem (1978  $\omega/KV$ , solved Alekseev-Meinrenken 2006  $\omega/AM$ , elucidated Alekseev-Torossian 2008-2012  $\omega/AT$ ), a problem about convolutions on Lie groups and Lie algebras.

**The Kashiwara-Vergne Conjecture.** There exist two series  $F$  and  $G$  in the completed free Lie algebra  $FL$  in generators  $x$  and  $y$  so that  $x+y - \log e^y e^x = (1 - e^{-\text{ad } x})F + (e^{\text{ad } y} - 1)G$  in  $FL$  and so that with  $z = \log e^x e^y$ ,

$$\text{tr}(\text{ad } x)\partial_x F + \text{tr}(\text{ad } y)\partial_y G \text{ in cyclic words} \\ = \frac{1}{2} \text{tr} \left( \frac{\text{ad } x}{e^{\text{ad } x} - 1} + \frac{\text{ad } y}{e^{\text{ad } y} - 1} - \frac{\text{ad } z}{e^{\text{ad } z} - 1} - 1 \right)$$

Implies the loosely-stated **convolutions statement**: Convolutions of invariant functions on a Lie group agree with convolutions of invariant functions on its Lie algebra.

**The Machine.** Let  $G$  be a group,  $\mathcal{K} = \mathbb{Q}G = \{\sum a_i g_i : a_i \in \mathbb{Q}, g_i \in G\}$  its group-ring,  $\mathcal{I} = \{\sum a_i g_i : \sum a_i = 0\} \subset \mathcal{K}$  its augmentation ideal. Let

$$\mathcal{A} = \text{gr } \mathcal{K} := \bigoplus_{m \geq 0} \mathcal{I}^m / \mathcal{I}^{m+1}.$$


**P.S.**  $(\mathcal{K}/\mathcal{I}^{m+1})^*$  is Vassiliev / finite-type / polynomial invariants.

Note that  $\mathcal{A}$  inherits a product from  $G$ .

**Definition.** A linear  $Z: \mathcal{K} \rightarrow \mathcal{A}$  is an "expansion" if for any  $\gamma \in \mathcal{I}^m$ ,  $Z(\gamma) = (0, \dots, 0, \gamma/\mathcal{I}^{m+1}, *, \dots)$ , and a "homomorphic expansion" if in addition it preserves the product.

**Example.** Let  $\mathcal{K} = C^\infty(\mathbb{R}^n)$  and  $\mathcal{I} = \{f : f(0) = 0\}$ . Then  $\mathcal{I}^m = \{f : f \text{ vanishes like } |x|^m\}$  so  $\mathcal{I}^m/\mathcal{I}^{m+1}$  degree  $m$  homogeneous polynomials and  $\mathcal{A} = \{\text{power series}\}$ . The Taylor series is a homomorphic expansion!


**Just for fun.**






In the finitely presented case, finding  $Z$  amounts to solving a system of equations in a graded space.

**Theorem** (with Zsuzsanna Dancso,  $\omega/WKO$ ). There is a bijection between the set of homomorphic expansions for  $w\mathcal{K}$  and the set of solutions of the Kashiwara-Vergne problem. **This is the tip of a major iceberg.**




Dancso,  $\omega/ZD$




4D Knots.




**The Generators**




**The Double Inflation Procedure.**



**wKO.**



**The Machine** generalizes to arbitrary algebraic structures!



"God created the knots, all else in topology is the work of mortals."

Leopold Kronecker (modified)

[www.katlas.org](http://www.katlas.org)





# A Partial Reduction of BF Theory to Combinatorics, 1

**Abstract.** I will describe a **semi-rigorous** reduction of perturbative BF theory (Cattaneo-Rossi [CR]) to computable combinatorics, in the case of ribbon 2-links. Also, I will explain how and why my approach may or may not work in the non-ribbon case. **Weak** this result is, and at least partially already known (Watanabe [Wa]). Yet in the ribbon case, the resulting invariant is a universal finite type invariant, a gadget that significantly generalizes and clarifies the Alexander polynomial and that is closely related to the Kashiwara-Vergne problem. I cannot rule out the possibility that the corresponding gadget in the non-ribbon case will be as interesting. (good news in **highlight**)

**The BF Feynman Rules.** For an edge  $e$ , let  $\Phi_e$  be its direction, in  $S^3$  or  $S^1$ . Let  $\omega_3$  and  $\omega_1$  be volume forms on  $S^3$  and  $S^1$ . Then for a 2-link  $(k_l)_{l \in T}$ ,



$$\zeta = \log \sum_{\text{diagrams } D} \frac{|D|}{|\text{Aut}(D)|} \int_{\mathbb{R}^2} \dots \int_{\mathbb{R}^2} \int_{\mathbb{R}^4} \dots \int_{\mathbb{R}^4} \prod_{\text{red } e \in D} \Phi_e^* \omega_3 \prod_{\text{black } e \in D} \Phi_e^* \omega_1$$






is an invariant in  $CW(FL(T)) \rightarrow CW(T)/\sim$ , "symmetrized cyclic words in  $T$ ".

**BF Following [CR].**  $A \in \Omega^1(M = \mathbb{R}^4, g)$ ,  $B \in \Omega^2(M, g^*)$ ,


$$S(A, B) := \int_M \langle B, F_A \rangle.$$

With  $\kappa: (S = \mathbb{R}^2) \rightarrow M$ ,  $\beta \in \Omega^0(S, g)$ ,  $\alpha \in \Omega^1(S, g^*)$ , set

$$O(A, B, \kappa) := \int \mathcal{D}\beta \mathcal{D}\alpha \exp\left(\frac{i}{\hbar} \int_S \langle \beta, d\kappa^* A \alpha + \kappa^* B \rangle\right).$$




**A 2-twist spun trefoil by Carter-Kamada-Saito [CKS].**




## A Partial Reduction of BF Theory to Combinatorics, 2

**Theorem 1 (with Cattaneo, Dalvit (credit, no blame)).** In the ribbon case,  $e^\zeta$  can be computed as follows:




**Sketch of Proof.** In 4D axial gauge, only “drop down” red propagators, hence in the ribbon case, no  $M$ -trivalent vertices.  $S$  integrals are  $\pm 1$  iff “ground pieces” run on nested curves as below, and exponentials arise when several propagators compete for the same double curve. And then the combinatorics is obvious...



### Musings

**Chern-Simons.** When the domain of BF is restricted to ribbon knots, and the target of Chern-Simons is restricted to trees and wheels, they agree. Why?

Is this all? What about the  $\vee$ -invariant?



**Theorem 2.** Using Gauss diagrams to represent knots and  $T$ -component pure tangles, the above formulas define an invariant in  $CW(FL(T)) \rightarrow CW(T)$ , “cyclic words in  $T$ ”.

- Agrees with BN-Dancso [BND] and with [BN2].
- In-practice computable!
- Vanishes on braids.
- Extends to w.
- Contains Alexander.
- The “missing factor” in Levine’s factorization [Le] (the rest of [Le] also fits, hence contains the MVA).
- Related to / extends Farber’s [Fa]?
- Should be summed and categorified.

### References.

[Ar] V. I. Arnold, *Topological Invariants of Plane Curves and Caustics*, University Lecture Series 5, American Mathematical Society 1994.

[BN1] D. Bar-Natan, *Bracelets and the Goussarov filtration of the space of knots, Invariants of knots and 3-manifolds (Kyoto 2001)*, Geometry and Topology Monographs 4 1–12, [arXiv:math.GT/0111267](http://arxiv.org/abs/math.GT/0111267).

[BN2] D. Bar-Natan, *Balloons and Hoops and their Universal Finite Type Invariant, BF Theory, and an Ultimate Alexander Invariant*, <http://www.math.toronto.edu/~drorbn/papers/KBH/>, [arXiv:1308.1721](http://arxiv.org/abs/1308.1721).

[BND] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects: From Alexander to Kashiwara and Vergne*, <http://www.math.toronto.edu/~drorbn/papers/WK0/>.

[CKS] J. S. Carter, S. Kamada, and M. Saito, *Diagrammatic Computations for Quandles and Cocycle Knot Invariants*, *Contemp. Math.* **318** (2003) 51–74.

[CS] J. S. Carter and M. Saito, *Knotted surfaces and their diagrams*, *Mathematical Surveys and Monographs* 55, American Mathematical Society, Providence 1998.

[Da] E. Dalvit, <http://science.unitn.it/~dalvit/>.

[CR] A. S. Cattaneo and C. A. Rossi, *Wilson Surfaces and Higher Dimensional Knot Invariants*, *Commun. in Math. Phys.* **256-3** (2005) 513–537, [arXiv:math-ph/0210037](http://arxiv.org/abs/math-ph/0210037).

[Fa] M. Farber, *Noncommutative Rational Functions and Boundary Links*, *Math. Ann.* **293** (1992) 543–568.

[Le] J. Levine, *A Factorization of the Conway Polynomial*, *Comment. Math. Helv.* **74** (1999) 27–53, [arXiv:q-alg/9711007](http://arxiv.org/abs/q-alg/9711007).


[Ro] D. Roseman, *Reidemeister-Type Moves for Surfaces in Four-Dimensional Space*, *Knot Theory*, Banach Center Publications 42 (1998) 347–380.

[Wa] T. Watanabe, *Configuration Space Integrals for Long  $n$ -Knots, the Alexander Polynomial and Knot Space Cohomology*, *Alg. and Geom. Top.* **7** (2007) 47–92, [arXiv:math/0609742](http://arxiv.org/abs/math/0609742).

(the “true” triple linking number)  $\bigcirc = \bigcup + \bigcap$


**Gnots.** In 3D, a generic immersion of  $S^1$  is an embedding, a knot. In 4D, a generic immersion of a surface has finitely-many double points (a gnot?). Perhaps we should be studying these?

**Finite type.** What are finite-type invariants for 2-knots? What would be “chord diagrams”?



**Bubble-wrap-finite-type.**

There’s an alternative definition of finite type in 3D, due to Goussarov (see [BN1]). The obvious parallel in 4D involves “bubble wraps”. Is it any good?




**Shielded tangles.** In 3D, one can’t zoom in and compute “the Chern-Simons invariant of a tangle”. Yet there are well-defined invariants of “shielded tangles”, and rules for their compositions. What would the 4D analog be?



Will the relationship with the Kashiwara-Vergne problem [BND] necessarily arise here?

**Plane curves.** Shouldn’t we understand integral / finite type invariants of plane curves, in the style of Arnold’s  $J^+$ ,  $J^-$ , and  $St$  [Ar], a bit better?




	$a(\times)$	$a(\times)$	$a(\times)$	$\infty$	$\bigcirc$	$\bigcirc$	$\bigcirc$	$\bigcirc$	$\dots$
St	1	0	0	0	0	1	2	3	$\dots$
$J^+$	0	2	0	0	0	-2	-4	-6	$\dots$
$J^-$	0	0	-2	-1	0	-3	-6	-9	$\dots$

### Continuing Joost Slingerland...

<http://youtu.be/YCA0VIEhVhg>

<http://youtu.be/mHyT0cf990>

“God created the knots, all else in topology is the work of mortals.”  
 Leopold Kronecker (modified)



**What are w-Trivalent Tangles?** (PA := Planar Algebra)


$\{ \text{knots} \} = \text{PA} \langle \text{R123} : \text{R123} \rangle$

$\{ \text{trivalent tangles} \} = \text{PA} \langle \text{R23, R4} \rangle$

wTT =  $\{ \text{trivalent w-tangles} \} = \text{PA} \langle \text{w-generators} \mid \text{w-relations} \mid \text{unary w-operations} \rangle$

**The w-generators.**

Broken surface, 2D Symbol, Dim. reduc., Virtual crossing, Movie



Cap, Wen, w, Vertices, smooth, singular


**Homomorphic expansions** for a filtered algebraic structure  $\mathcal{K}$ :

$$\text{ops}^{\leftarrow} \mathcal{K} = \mathcal{K}_0 \supset \mathcal{K}_1 \supset \mathcal{K}_2 \supset \mathcal{K}_3 \supset \dots$$

$$\text{ops}^{\leftarrow} \text{gr } \mathcal{K} := \mathcal{K}_0/\mathcal{K}_1 \oplus \mathcal{K}_1/\mathcal{K}_2 \oplus \mathcal{K}_2/\mathcal{K}_3 \oplus \mathcal{K}_3/\mathcal{K}_4 \oplus \dots$$

An **expansion** is a filtration  $Z : \mathcal{K} \rightarrow \text{gr } \mathcal{K}$  that "covers" the identity on  $\text{gr } \mathcal{K}$ . A **homomorphic expansion** is an expansion that respects all relevant "extra" operations.

**Filtered algebraic structures are cheap and plenty.** In any  $\mathcal{K}$ , allow formal linear combinations, let  $\mathcal{K}_1$  be the ideal generated by differences (the "augmentation ideal"), and let  $\mathcal{K}_m := \langle (\mathcal{K}_1)^m \rangle$  (using all available "products").



- Has kinds, objects, operations, and maybe constants.
- Perhaps subject to some axioms.
- We always allow formal linear combinations.

**Example: Pure Braids.**  $PB_n$  is generated by  $x_{ij}$ , "strand  $i$  goes around strand  $j$  once", modulo "Reidemeister moves".  $A_n := \text{gr } PB_n$  is generated by  $t_{ij} := x_{ij} - 1$ , modulo the 4T relations  $[t_{ij}, t_{ik} + t_{jk}] = 0$  (and some lesser ones too). Much happens in  $A_n$ , including the Drinfel'd theory of associators.

**Our case(s).**


$\mathcal{K} \xrightarrow{Z: \text{high algebra}} \mathcal{A} := \text{gr } \mathcal{K} \xrightarrow{\text{given a "Lie" algebra } \mathfrak{g}} \mathcal{U}(\mathfrak{g})$

solving finitely many equations in finitely many unknowns

low algebra: pictures represent formulas

$\mathcal{K}$  is knot theory or topology;  $\text{gr } \mathcal{K}$  is finite combinatorics: bounded-complexity diagrams modulo simple relations.

A **Ribbon 2-Knot** is a surface  $S$  embedded in  $\mathbb{R}^4$  that bounds an immersed handlebody  $B$ , with only "ribbon singularities"; a ribbon singularity is a disk  $D$  of transverse double points, whose preimages in  $B$  are a disk  $D_1$  in the interior of  $B$  and a disk  $D_2$  with  $D_2 \cap \partial B = \partial D_2$ , modulo isotopies of  $S$  alone.



The **w-relations** include R234, VR1234, M, Overcrossings Commute (OC) but not UC,  $W^2 = 1$ , and funny interactions between the wen and the cap and over- and under-crossings:

OC:  $\text{OC} : \text{crossing} \leftrightarrow \text{crossing}$  as  $\text{crossing} \leftrightarrow \text{crossing}$  yet not UC:  $\text{UC} : \text{crossing} \leftrightarrow \text{crossing}$

Challenge. Do the Reidemeister!

The unary w-operations

Unzip along an annulus, Unzip along a disk

**Just for fun.**

$\mathcal{K} = \{ \text{Reidemeister}, \text{Winter} \} = \left( \text{The set of all b/w 2D projections of reality} \right)$

$\mathcal{K}/\mathcal{K}_1 \leftarrow \mathcal{K}/\mathcal{K}_2 \leftarrow \mathcal{K}/\mathcal{K}_3 \leftarrow \mathcal{K}/\mathcal{K}_4 \leftarrow \dots$

Crop Rotate Adjoin


An expansion  $Z$  is a choice of a "progressive scan" algorithm.

$\mathcal{K}/\mathcal{K}_1 \oplus \mathcal{K}_1/\mathcal{K}_2 \oplus \mathcal{K}_2/\mathcal{K}_3 \oplus \mathcal{K}_3/\mathcal{K}_4 \oplus \mathcal{K}_4/\mathcal{K}_5 \oplus \mathcal{K}_5/\mathcal{K}_6 \oplus \dots$


$\mathbb{R} \parallel \ker(\mathcal{K}/\mathcal{K}_4 \rightarrow \mathcal{K}/\mathcal{K}_3)$

# Convolutions on Lie Groups and Lie Algebras and Ribbon 2-Knots, Page 2

**Knot-Theoretic statement.** There exists a homomorphic expansion  $Z$  for trivalent w-tangles. In particular,  $Z$  should respect  $R4$  and intertwine annulus and disk unzips:



**Diagrammatic statement.** Let  $R = \exp \uparrow \in \mathcal{A}^w(\uparrow\uparrow)$ . There exist  $\omega \in \mathcal{A}^w(\uparrow)$  and  $V \in \mathcal{A}^w(\uparrow\uparrow)$  so that



**Algebraic statement.** With  $I\mathfrak{g} := \mathfrak{g}^* \rtimes \mathfrak{g}$ , with  $c : \hat{U}(I\mathfrak{g}) \rightarrow \hat{U}(\mathfrak{g})/\hat{U}(\mathfrak{g}) = \hat{S}(\mathfrak{g}^*)$  the obvious projection, with  $S$  the antipode of  $\hat{U}(I\mathfrak{g})$ , with  $W$  the automorphism of  $\hat{U}(I\mathfrak{g})$  induced by flipping the sign of  $\mathfrak{g}^*$ , with  $r \in \mathfrak{g}^* \otimes \mathfrak{g}$  the identity element and with  $R = e^r \in \hat{U}(I\mathfrak{g}) \otimes \hat{U}(\mathfrak{g})$  there exist  $\omega \in \hat{S}(\mathfrak{g}^*)$  and  $V \in \hat{U}(I\mathfrak{g})^{\otimes 2}$  so that

(1)  $V(\Delta \otimes 1)(R) = R^{13}R^{23}V$  in  $\hat{U}(I\mathfrak{g})^{\otimes 2} \otimes \hat{U}(\mathfrak{g})$   
 (2)  $V \cdot SWV = 1$       (3)  $(c \otimes c)(V\Delta(\omega)) = \omega \otimes \omega$

**Unitary statement.** There exists  $\omega \in \text{Fun}(\mathfrak{g})^G$  and an (infinite order) tangential differential operator  $V$  defined on  $\text{Fun}(\mathfrak{g}_x \times \mathfrak{g}_y)$  so that

(1)  $V\widehat{e^{x+y}} = \widehat{e^x e^y} V$  (allowing  $\hat{U}(\mathfrak{g})$ -valued functions)  
 (2)  $VV^* = I$       (3)  $V\omega_{x+y} = \omega_x \omega_y$


**Group-Algebra statement.** There exists  $\omega^2 \in \text{Fun}(\mathfrak{g})^G$  so that for every  $\phi, \psi \in \text{Fun}(\mathfrak{g})^G$  (with small support), the following holds in  $\hat{U}(\mathfrak{g})$ :

$$\iint_{\mathfrak{g} \times \mathfrak{g}} \phi(x)\psi(y)\omega_{x+y}^2 e^{x+y} = \iint_{\mathfrak{g} \times \mathfrak{g}} \phi(x)\psi(y)\omega_x^2 \omega_y^2 e^x e^y. \quad (\text{shhh, this is Duflo})$$


**Convolutions statement** (Kashiwara-Vergne). Convolutions of invariant functions on a Lie group agree with convolutions of invariant functions on its Lie algebra. More accurately, let  $G$  be a finite dimensional Lie group and let  $\mathfrak{g}$  be its Lie algebra, let  $j : \mathfrak{g} \rightarrow \mathbb{R}$  be the Jacobian of the exponential map  $\exp : \mathfrak{g} \rightarrow G$ , and let  $\Phi : \text{Fun}(G) \rightarrow \text{Fun}(\mathfrak{g})$  be given by  $\Phi(f)(x) := j^{1/2}(x)f(\exp x)$ . Then if  $f, g \in \text{Fun}(G)$  are Ad-invariant and supported near the identity, then

$$\Phi(f) \star \Phi(g) = \Phi(f \star g).$$


**From wTT to  $\mathcal{A}^w$ .**  $\text{gr}_m \text{wTT} := \{m\text{-cubes}\} / \{(m+1)\text{-cubes}\}$ :



**w-Jacobi diagrams and  $\mathcal{A}$ .**  $\mathcal{A}^w(Y \uparrow) \cong \mathcal{A}^w(\uparrow\uparrow\uparrow)$  is



**Diagrammatic to Algebraic.** With  $(x_i)$  and  $(\varphi^j)$  dual bases of  $\mathfrak{g}$  and  $\mathfrak{g}^*$  and with  $[x_i, x_j] = \sum b_{ij}^k x_k$ , we have  $\mathcal{A}^w \rightarrow \mathcal{U}$  via




**Unitary  $\iff$  Algebraic.** The key is to interpret  $\hat{U}(I\mathfrak{g})$  as tangential differential operators on  $\text{Fun}(\mathfrak{g})$ :

- $\varphi \in \mathfrak{g}^*$  becomes a multiplication operator.
- $x \in \mathfrak{g}$  becomes a tangential derivation, in the direction of the action of  $\text{ad } x$ :  $(x\varphi)(y) := \varphi([x, y])$ .
- $c : \hat{U}(I\mathfrak{g}) \rightarrow \hat{U}(I\mathfrak{g})/\hat{U}(\mathfrak{g}) = \hat{S}(\mathfrak{g}^*)$  is “the constant term”.

**Unitary  $\implies$  Group-Algebra.**  $\iint \omega_{x+y}^2 e^{x+y} \phi(x)\psi(y) = \langle \omega_{x+y}, \omega_{x+y} e^{x+y} \phi(x)\psi(y) \rangle = \langle V\omega_{x+y}, V e^{x+y} \phi(x)\psi(y)\omega_{x+y} \rangle = \langle \omega_x \omega_y, e^x e^y V \phi(x)\psi(y)\omega_{x+y} \rangle = \langle \omega_x \omega_y, e^x e^y \phi(x)\psi(y)\omega_x \omega_y \rangle = \iint \omega_x^2 \omega_y^2 e^x e^y \phi(x)\psi(y).$

**Convolutions and Group Algebras** (ignoring all Jacobians). If  $G$  is finite,  $A$  is an algebra,  $\tau : G \rightarrow A$  is multiplicative then  $(\text{Fun}(G), \star) \cong (A, \cdot)$  via  $L : f \mapsto \sum f(a)\tau(a)$ . For Lie  $(G, \mathfrak{g})$ ,



with  $L_0\psi = \int \psi(x)e^x dx \in \hat{S}(\mathfrak{g})$  and  $L_1\Phi^{-1}\psi = \int \psi(x)e^x \in \hat{U}(\mathfrak{g})$ . Given  $\psi_i \in \text{Fun}(\mathfrak{g})$  compare  $\Phi^{-1}(\psi_1) \star \Phi^{-1}(\psi_2)$  and  $\Phi^{-1}(\psi_1 \star \psi_2)$  in  $\hat{U}(\mathfrak{g})$ : (shhh,  $L_{0/1}$  are “Laplace transforms”)

$\star$  in  $G$ :  $\iint \psi_1(x)\psi_2(y)e^x e^y$        $\star$  in  $\mathfrak{g}$ :  $\iint \psi_1(x)\psi_2(y)e^{x+y}$

- We skipped...**
- The Alexander polynomial and Milnor numbers.
  - v-Knots, quantum groups and Etingof-Kazhdan.
  - u-Knots, Alekseev-Torossian, and BF theory and the successful and Drinfel'd associators.
  - The simplest problem hyperbolic geometry solves.
  - The religion of path integrals.