



**Abstract.** I will describe a **computable, non-commutative** invariant of tangles with values in wheels, almost generalize it to some balloons, and then tell you why I care. Spoilers: tangles are you know what, wheels are linear combinations of cyclic words in some alphabet, balloons are 2-knots, and one reason I care is because quantum field theory predicts more than I can actually get (but also less).

**Why I like "non-commutative"?** With  $FA(x_i)$  the free associative non-commutative algebra,


$$\dim \mathbb{Q}[x, y]_d \sim d \ll 2^d \sim \dim FA(x, y)_d.$$

**Why I like "computable"?**


- Because I'm weird.
- Note that  $\pi_1$  isn't computable.

**Preliminaries from Algebra.**  $FL(x_i)$

denotes the free Lie algebra in  $(x_i)$ ;  $FL(x_i) = (\text{binary trees with AS vertices and coloured leaves}) / (\text{IHX relations})$ . There an obvious map  $FA(FL(x_i)) \rightarrow FA(x_i)$  defined by  $[a, b] \rightarrow ab - ba$ , which in itself, is IHX.




$CW(x_i)$  denotes the vector space of cyclic words in  $(x_i)$ :  $CW(x_i) = FA(x_i) / (x_i w = w x_i)$ . There an obvious map  $CW(FL(x_i)) \rightarrow CW(x_i)$ . In fact, connected uni-trivalent 2-in-1-out graphs with univalents with colours in  $\{1, \dots, n\}$ , modulo AS and IHX, is precisely  $CW(x_i)$ :





**Most important.**  $e^x = \sum \frac{x^d}{d!}$  and  $e^{x+y} = e^x e^y$ .


**Preliminaries from Knot Theory.**




**Theorem.**  $\omega$ , the connected part of the procedure below, is an invariant of  $S$ -component tangles with values in  $CW(S)$ :



reconsider colour coding



$\omega$  is practically computable! For the Borromean tangle, to degree 5, the result is: (see [BN])




**Proof of Invariance.**

Need to show:

$$\omega \left( \begin{array}{c} \uparrow \uparrow \uparrow \\ \leftarrow \rightarrow \leftarrow \rightarrow \\ \uparrow \uparrow \uparrow \end{array} \right) = \omega \left( \begin{array}{c} \uparrow \uparrow \uparrow \\ \leftarrow \rightarrow \leftarrow \rightarrow \\ \uparrow \uparrow \uparrow \end{array} \right)$$

Indeed,





(thanks, Ester Dalvit)

- $\omega$  is really the second part of a (trees, wheels)-valued **Further Facts** invariant  $\zeta = (\lambda, \omega)$ . The tree part  $\lambda$  is just a repackaging of the Milnor  $\mu$ -invariants.
- On u-tangles,  $\zeta$  is equivalent to the trees&wheels part of the Kontsevich integral, except it is computable and is defined with no need for a choice of parenthesization.
- On long/round u-knots,  $\omega$  is equivalent to the Alexander polynomial.
- The multivariable Alexander polynomial (and Levine's factorization thereof [Le]) is contained in the Abelianization of  $\zeta$  [BNS].
- $\omega$  vanishes on braids.
- Related to / extends Farber's [Fa]?
- Should be summed and categorified.
- Extends to v and descends to w: meaning,  $\zeta$  satisfies  $\omega$  also satisfies


meaning,  $\zeta$  satisfies

$\omega$  also satisfies




- Agrees with BN-Dancso [BND1, BND2] and with [BN].
- Using  $\mathcal{K}: v\mathcal{K}_n \rightarrow w\mathcal{K}_{n+1}$ , defines a strong invariant of v-tangles / long v-knots.

$\notin \text{LaTeX}$ : web/the



2-Knots.



**A Big Question.** Does it all extend to arbitrary 2-knots (not necessarily “simple”)? To arbitrary codimension-2 knots?

**BF Following [CR].**  $A \in \Omega^1(M = \mathbb{R}^4, \mathfrak{g}), B \in \Omega^2(M, \mathfrak{g}^*),$

$$S(A, B) := \int_M \langle B, F_A \rangle.$$

With  $\kappa: (S = \mathbb{R}^2) \rightarrow M, \beta \in \Omega^0(S, \mathfrak{g}), \alpha \in \Omega^1(S, \mathfrak{g}^*),$  set

$$\mathcal{O}(A, B, \kappa) := \int \mathcal{D}\beta \mathcal{D}\alpha \exp\left(\frac{i}{\hbar} \int_S \langle \beta, d_{\kappa^*} \alpha + \kappa^* B \rangle\right).$$



Rossi


**The BF Feynman Rules.** For an edge  $e,$  let  $\Phi_e$  be its direction, in  $S^3$  or  $S^1.$  Let  $\omega_3$  and  $\omega_1$  be volume forms on  $S^3$  and  $S^1.$  Then

$$Z_{BF} = \sum_{\text{diagrams } D} \frac{|D|}{|\text{Aut}(D)|} \underbrace{\int_{\mathbb{R}^2} \dots \int_{\mathbb{R}^2}}_{S\text{-vertices}} \underbrace{\int_{\mathbb{R}^4} \dots \int_{\mathbb{R}^4}}_{M\text{-vertices}} \prod_{e \in D} \Phi_e^* \omega_3 \prod_{e \in D} \Phi_e^* \omega_1$$



Cattaneo


(modulo some IHX-like relations).




**Issues.** • Signs don't quite work out, and BF seems to reproduce only “half” of the wheels invariant.

- There are many more configuration space integrals than BF Feynman diagrams and than just trees and wheels.
- I don't know how to define “finite type” for arbitrary 2-knots.

The Generators




The Double Inflation Procedure  $\delta.$



The Full

2-Knot Story



- Missing:
1. The Roseman moves?
  2. Satoh's conjecture?
  3. A picture of Balloons?  $\circ\circ\circ$

References.

[BN] D. Bar-Natan, *Balloons and Hoops and their Universal Finite Type 1-Invariant, BF Theory, and an Ultimate Alexander Invariant*,  $\omega\epsilon\beta/\text{KBH}$ , arXiv:1308.1721.

[BND1] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects I: W-Knots and the Alexander Polynomial*,  $\omega\epsilon\beta/\text{WKO1}$ , arXiv:1405.1956.

[BND2] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects II: Tangles and the Kashiwara-Vergne Problem*,  $\omega\epsilon\beta/\text{WKO2}$ , arXiv:1405.1955.

[BNS] D. Bar-Natan and S. Selmani, *Meta-Monoids, Meta-Bicrossed Products, and the Alexander Polynomial*, J. of Knot Theory and its Ramifications **22-10** (2013), arXiv:1302.5689.

[CS] J. S. Carter and M. Saito, *Knotted surfaces and their diagrams*, Math. Surv. and Mono. **55**, Amer. Math. Soc., Providence 1998.

[CR] A. S. Cattaneo and C. A. Rossi, *Wilson Surfaces and Higher Dimensional Knot Invariants*, Commun. in Math. Phys. **256-3** (2005) 513–537, arXiv:math-ph/0210037.

[Fa] M. Farber, *Noncommutative Rational Functions and Boundary Links*, Math. Ann. **293** (1992) 543–568.

[Le] J. Levine, *A Factorization of the Conway Polynomial*, Comment. Math. Helv. **74** (1999) 27–53, arXiv:q-alg/9711007.

[Wa] T. Watanabe, *Configuration Space Integrals for Long n-Knots, the Alexander Polynomial and Knot Space Cohomology*, Alg. and Geom. Top. **7** (2007) 47–92, arXiv:math/0609742.



“God created the knots, all else in topology is the work of mortals.”

Leopold Kronecker (modified)

