

Name: Dror Bar-Natan

Student ID: Grading Key

UNIVERSITY OF TORONTO  
Faculty of Arts and Sciences  
DECEMBER EXAMINATIONS 2014  
Math 240H1 Algebra I — Final Exam

Dror Bar-Natan  
December 10, 2014

Solve 4 of the following 5 questions. If you solve more than 4 questions indicate very clearly below which are the ones that you want marked, or else a random one will be left out. Please write your answers within this booklet. You may also use the back side of the pages for answers and you may continue answers in the blank pages provided at the end of this notebook (though indicate "continued on page ..." when you do so). You may mark pages "draft" and use them for scratch work.

The questions carry equal weight though different parts of the same question may be weighted differently.

There are 3 pages in this exam booklet.

**Duration.** You have 3 hours to write this exam.

**Allowed Material.** Stationary and basic calculators, not capable of displaying text or sounding speech. A stuffed animal for personal comfort.

**Good Luck!**

For grading use — circle the 4 questions that you want marked:

1	/25	4	/25
2	/25	5	/25
3	/25		
Total	/100		

**Problem 1.** Let  $A$  be a matrix in  $M_{n \times n}(F)$ . Let  $T: M_{n \times n}(F) \rightarrow M_{n \times n}(F)$  be defined by  $T(B) = AB$ .

1. Show that  $T$  is a linear transformation.

2. Find the matrix representing  $T$  relative to the basis

$$\left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right\},$$

which is taken as a basis of both the domain space of  $T$  and the target space of  $T$ .

**Tip.** In math-talk, "show" means "prove".

**Tip.** Don't start working! Read the whole exam first. You may wish to start with the questions that are easiest for you.

**Tip.** Neatness, cleanliness and organization count, here and everywhere else!

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \rightarrow \begin{pmatrix} a & 0 & b & 0 \\ 0 & a & 0 & b \\ c & 0 & d & 0 \\ 0 & c & 0 & d \end{pmatrix} = \begin{pmatrix} a_{11} & 0 & a_{12} & 0 \\ 0 & a_{11} & 0 & a_{12} \\ a_{21} & 0 & a_{22} & 0 \\ 0 & a_{21} & 0 & a_{22} \end{pmatrix}$$

-4 Chose a specific matrix  $A$   $\left( \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \right)$ .

-7  $T \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} a_{11} & 0 \\ 0 & 0 \end{pmatrix}$  etc.

-5 computed  $M$  assuming mult. on the right.

FTTFF | TFFT | ~~ET~~ TT  
TF

**Problem 2.** State and prove the "dimension theorem", also known as the "rank-nullity theorem", for a given linear transformation  $T: V \rightarrow W$ .

**Tip.** In math-talk, "state" means "write the statement of, in full".

**Tip.** As always in math exams, when proving a theorem you may freely assume anything that preceded it but you may not assume anything that followed it.

state: 10  
prove: 15

---

state.

---

prove. 2/15 assumed  $T$  is invertible then did well.

(-7) completely wrong pt. of lin. indep.  
of  $\{T(v_i)\}$ .

(-4) "Since  $u_1, \dots, u_n$  exclude the basis for  $N(T)$ ". . .

FTTFF | TFFT | TFFT

**Problem 3.** Find all the solutions (if any exist) of the following two systems of linear equations:

$$\begin{pmatrix} 0 & 0 & 1 & 3 \\ 1 & -2 & 0 & -1 \\ 1 & -2 & 1 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 2 \\ 3 \\ 3 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 0 & 0 & 1 & 3 \\ 1 & -2 & 0 & -1 \\ 1 & -2 & 1 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}.$$

**Tip.** Show all intermediate steps!

**Tip.** It is always an excellent idea to substitute your solutions back into the equations and see if they really work.

$$\begin{pmatrix} 0 & 0 & 1 & 3 & | & 2 & 1 \\ 1 & -2 & 0 & -1 & | & 3 & 1 \\ 1 & -2 & 1 & 2 & | & 3 & 2 \end{pmatrix} \mapsto \begin{pmatrix} 1 & -2 & 0 & -1 & | & 3 & 1 \\ 0 & 0 & 1 & 3 & | & 0 & 1 \\ 0 & 0 & 1 & 3 & | & 2 & 1 \end{pmatrix} \mapsto \begin{pmatrix} 1 & -2 & 0 & -1 & | & 3 & 1 \\ 0 & 0 & 1 & 3 & | & 0 & 1 \\ 0 & 0 & 0 & 0 & | & 2 & 0 \end{pmatrix}$$

Sys 1: no solns.

Sys 2:  $x_1 - 2x_2 - x_4 = 1$     soln:  $\begin{pmatrix} 1 + 2x_2 + x_4 \\ x_2 \\ 1 - 3x_4 \\ x_4 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix} + \begin{pmatrix} 2 \\ 1 \\ 0 \\ 0 \end{pmatrix} x_2 + \begin{pmatrix} 1 \\ 0 \\ -3 \\ 1 \end{pmatrix} x_4$   
 $x_3 + 3x_4 = 1$

~~10~~ In part II, correct row reduction but completely wrong "reading of answer". No check.

~~6~~ ~~10~~ Forgot the inhomogeneous part  $\begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix}$ , no check.

~~4~~ ~~4~~ Wrong end result, no checks.

~~8~~ Found just one soln.

F T T F F | T F F T T | T F T T

Problem 4. Let  $A$  be the matrix  $A = \begin{pmatrix} 5 & -3 \\ 6 & -4 \end{pmatrix}$ .

1. Compute  $\det(A - \lambda I)$ .  $(5-\lambda)(-4-\lambda) - (-3) \cdot 6 = \lambda^2 - \lambda - 2$
2. Find the eigenvalues  $\lambda_1$  and  $\lambda_2$  of  $A$ .  $\lambda_{1,2} = -1, 2$
3. Find their corresponding eigenvectors  $v_1$  and  $v_2$ .  $\begin{pmatrix} 5-\lambda_1 & -3 \\ 6 & -4+\lambda_1 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = 0 \Rightarrow v_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$
4. Find a matrix  $C$  for which  $AC = CD$ , where  $D = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}$ .  $\begin{pmatrix} 3 & -3 \\ 6 & -6 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = 0 \Rightarrow v_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$
5. Compute the inverse of  $C$ .  $C^{-1} = \begin{pmatrix} -1 & 1 \\ 2 & -1 \end{pmatrix}$ ,  $C = \begin{pmatrix} 1 & 1 \\ 2 & 1 \end{pmatrix}$
6. For an arbitrary natural number  $n$ , compute  $A^n$  by computing  $CD^nC^{-1}$ .

$$A^n = \begin{pmatrix} 1 & 1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} (-1)^n & 0 \\ 0 & 2^n \end{pmatrix} \begin{pmatrix} -1 & 1 \\ 2 & -1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} -(-1)^n & (-1)^n \\ 2 \cdot 2^n & -2^n \end{pmatrix} = \begin{pmatrix} 2^{n+1} - (-1)^n & (-1)^n - 2^n \\ 2^{n+1} - 2(-1)^n & 2(-1)^n - 2^n \end{pmatrix}$$

**Problem 5.** In the spaces provided, indicate for each of the following statements if it is TRUE or FALSE. You may leave an entry blank. You will earn 2 points for each correct answer, (-3) points for each incorrect answer, and 0 points for each entry left blank. Your overall mark for this question will be raised to 0 if you earn a negative number of points.

If nothing is otherwise stated,  $A$  and  $B$  are assumed to be square  $n \times n$  matrices with entries in some field  $F$ .

1.   F   The function  $\det: M_{n \times n}(F) \rightarrow F$  is a linear transformation.
2.   T   The determinant of  $A$  can be evaluated using an expansion along any row or column.
3.   T   If two rows of  $A$  are identical then  $\det(A) = 0$ .
4.   F   If a row of  $A$  is identical to one of the columns of  $A$ , then  $\det(A) = 0$ .
5.   F   If  $AB \neq BA$  then  $\det(AB) \neq \det(BA)$ .
6.   T   If  $B$  is obtained from  $A$  by interchanging any two columns, then  $\det(B) = -\det(A)$ .
7.   F   If  $B$  is obtained from a  $A$  by multiplying a row of  $A$  by a scalar, then  $\det(B) = \det(A)$ .
8.   F   If  $B$  is obtained from  $A$  by adding  $c$  times one row to another row, then  $\det(B) = c \det(A)$ .
9.   T   If  $A$  has rank  $n$ , then  $\det(A) \neq 0$ .
10.   T   If  $A$  is upper triangular (meaning that  $A_{ij} = 0$  whenever  $i > j$ ), then  $\det(A)$  is equal to the product of the diagonal entries  $A_{ii}$  of  $A$ .
11.   T   Always,  $\det(A) = \det(A^T)$ .
12.   F   If  $A$  is invertible, then  $\det(A) = \det(A^{-1})$ .
13.   T   If  $A$  is an  $n \times n$  matrix where  $n$  is even, then  $\det(A) = \det(-A)$ .
14.   T   Always,  $\det(AB) = \det(BA)$ .

**Tip.** There is no need to justify your answers.

**Tip.** Note that a statement is considered TRUE only if it is **always** true, and not just **sometimes**.