


See a paper-folding solution at <http://www.make-origami.com/HelenaVerrill/perimeter.php>



Old riddles (sol'n at end). Can you fold a rectangular piece of paper (perhaps many times) so that the result will have a longer perimeter than the original? If  $f: A \rightarrow \mathbb{R}^2$  is a distance-non-increasing map from a rectangle to the plane, is it always the case that the length of the boundary of  $f(A)$  is less than the length of the boundary of  $A$ ?

Blame COVID!



Next tasks: IF  $M^3 \subset \mathbb{R}^3$  and  $F$  is a vector field,

$$\int_M \operatorname{div} F dV = \int_{\partial M} F \cdot n dA$$

*n* unit normal to  $\partial M$

IF  $M^2 \subset \mathbb{R}^3$  is compact orient,

$$\int_M (\operatorname{curl} F) \cdot n dA = \int_{\partial M} F \cdot T ds$$

*n* unit normal to  $M$ , *T* unit tangent to  $\partial M$


Integration on on ?

From Munkres' Analysis on Manifolds:

**Theorem 35.2.** Let  $M$  be a compact oriented  $k$ -manifold in  $\mathbb{R}^n$ . Let  $\omega$  be a  $k$ -form defined in an open set of  $\mathbb{R}^n$  containing  $M$ . Suppose that  $\alpha_i: A_i \rightarrow M_i$ , for  $i = 1, \dots, N$ , is a coordinate patch on  $M$  belonging to the orientation of  $M$ , such that  $A_i$  is open in  $\mathbb{R}^k$  and  $M$  is the disjoint union of the open sets  $M_1, \dots, M_N$  of  $M$  and a set  $K$  of measure zero in  $M$ . Then

$$\int_M \omega = \sum_{i=1}^N \left[ \int_{A_i} \alpha_i^* \omega \right]$$

But first, what are  $dV$ ,  $dA$ ,  $ds$ ?



by some  $\eta \in \mathcal{L}^k(M)$  defined up to mult by  $\mathbb{R}$ .


$M^k \subset \mathbb{R}^n$  oriented the volume form  $dV$  on  $M^k$  ( $dV \in \mathcal{L}^k(M)$ )

Warning  $\nabla$  isn't a function,  $dV$  isn't a 1-form,

by declaring that  $dV$  is that multiple of  $\eta$  (by a function) for which


$$dV(u_1, \dots, u_k) = 1$$

if  $u_1, \dots, u_k$  make an orthonormal basis of  $T_x M$  which agrees w/ the orientation of  $M$ .



↑ This is indep of the choice of O.N. basis b/c change of basis matrices between O.N. bases

Examples ①




have  $\det = \pm 1$

$$dV(T) = dS(T) = 1$$

$$\textcircled{2} S^2 = \{(x, y, z) : x^2 + y^2 + z^2 = 1\}$$

$$P = (0, 0, 1) \quad u = (1, 0, 0)$$

$$v = (0, 1, 0)$$



$$\langle dx \wedge dy \rangle(u, v) = \alpha \implies \alpha = 1$$

$$dA_p = dx \wedge dy$$



In general, if  $M^2 \subset \mathbb{R}^3$  is an oriented 2-manifold in  $\mathbb{R}^3$ , &  $n : M \rightarrow \bigcup_{x \in M} T_x \mathbb{R}^3$

s.t.,

1.  $n(x) \in T_x \mathbb{R}^3$
2.  $n(x) \perp T_x M, |n(x)| = 1$

"a unit normal to M"

3. If  $u, v$  define the orientation of  $M$  at  $x$ , then  $(u, v, n)$  define the orientation of  $\mathbb{R}^3$  at  $x$ .

In this case,


$$dV^{(u,v)} = dA(u, v) = \begin{vmatrix} -u- \\ -v- \\ -n- \end{vmatrix} = (u \times v) \cdot n$$

any two  $u, v \in T_x M$

bilinear & alternating as function of  $u, v$ .  
& has the right value (1) if  $(u, v)$  make an oriented ON. basis.

$$\begin{pmatrix} u_2 v_3 - u_3 v_2 \\ u_3 v_1 - u_1 v_3 \\ u_1 v_2 - u_2 v_1 \end{pmatrix} \cdot \begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix}$$

Reminder  $u \times v$  is normal to both  $u, v$ ,  $|u \times v| = \text{area of the parallelogram spanned by } u, v$ .  
 $(u, v, u \times v)$  defined the orientation of  $\mathbb{R}^3$




$$dA(u, v) = \begin{vmatrix} -u- \\ -v- \\ -n- \end{vmatrix} = (u \times v) \cdot n$$

$$= (n_1 dy_1 dz + n_2 dz_1 dx + n_3 dx_1 dy)(u, v)$$

$$(dy_1 dz)(u, v) = u_2 v_3 - v_2 u_3 \dots$$

Example What's  $dA$  on  $S^2$  at  $\begin{pmatrix} x \\ y \\ z \end{pmatrix} \in S^2$



$$p = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \quad n = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

$$dA = n_1 dy dz + \dots$$

$$= x dy dz + y dz dx + z dx dy$$

(as seen earlier)

$$\left(\frac{1}{1-x}\right)' \Big|_{x=\frac{1}{2}} = (1+x+x^2+\dots)' \Big|_{x=\frac{1}{2}} = (0+1x^0+2x^1+3x^2+\dots) \Big|_{x=\frac{1}{2}}$$


$$= \underbrace{1\left(\frac{1}{2}\right)^0}_{2} + 2\left(\frac{1}{2}\right)^1 + \underbrace{3\left(\frac{1}{2}\right)^2 + 4\left(\frac{1}{2}\right)^3 + \dots}_{\text{what we want}}$$

$$w_i = 1 + \frac{1}{2} + \frac{1}{2} = 2$$

$$w_e \leftarrow \sum_{k=2}^{\infty} 2^{-k} (k+1) = \left(\frac{1}{1-x}\right)' \Big|_{x=\frac{1}{2}} - 2$$

$$= \frac{1}{(1-x)^2} \Big|_{x=\frac{1}{2}} - 2 = 2$$


A move:




8								
7								
6	X	X						
5		X	X	X				
4	1/8		X	X	X			
3	1/4	<del>1/8</del>	X	X	X			
2	<del>1/2</del>	<del>1/4</del>	1/8	X	X	X		
1	<del>1</del>	<del>1/2</del>	1/4	1/8				
	A	B	C	D	E	F	G	H

Read Along: Spivak 126 to infinity.

Old riddle (sol'n at end). On a chessboard, there are three pawns at the lower left (at A1, A2, and B1). On each move, pick up one pawn, remove it and place one new pawn to the right and one new pawn above, but only if these squares are unoccupied. Can you clear the original 3 pawns?



A move:



Next tasks: IF  $M^3 \subset \mathbb{R}^3$   
and  $F$  is a vector field,

$$\int_M \operatorname{div} F dV = \int_{\partial M} F \cdot n dA$$

$\uparrow$  unit normal to  $\partial M$

IF  $M^2 \subset \mathbb{R}^3$  is compact orient,

$$\int_M (\operatorname{curl} F) \cdot n dA = \int_{\partial M} F \cdot T ds$$

$\uparrow$  unit normal to  $M$        $\uparrow$  unit tangent to  $\partial M$

But first, what are  $dV$ ,  
 $dA$ ,  $ds$ ?


IF  $M^2 \subset \mathbb{R}^3$  &  $n$  is the positive unit normal,

$$dA(u, v) = \begin{vmatrix} -u \\ -v \\ -n \end{vmatrix} = (u \times v) \cdot n = (n_1 dy dz + n_2 dz dx + n_3 dx dy)(u, v)$$

$$= \pm |u \times v| = \pm |u| |v| \sin \alpha = \pm \sqrt{|u|^2 |v|^2 - (u \cdot v)^2}$$

$\uparrow$  + if  $(u, v)$        $\uparrow$   $\sqrt{1 - \cos^2 \alpha}$        $\cos \alpha = \frac{u \cdot v}{|u| |v|}$

make a positive basis (basis that agrees w/ orientation)




Example Suppose  $M$  is the image of a single cube:  $M = C(\mathbb{I}^2)$  where  $C$  is smooth, 1-1, w/ injective  $C'$ .

$$\operatorname{Area}(M) = \int_M dA$$

$$\int_{I^2} C^*(dA) = \int_{I^2} \underbrace{C^*(dA)(e_1, e_2)} dx dy$$


$$= \int_{I^2} dA(C_* e_1, C_* e_2) = \int_{I^2} dA \left( \underbrace{\frac{\partial C}{\partial x}}_u, \underbrace{\frac{\partial C}{\partial y}}_v \right)$$

$$\begin{pmatrix} \frac{\partial C_1}{\partial x} & \frac{\partial C_1}{\partial y} \\ \frac{\partial C_2}{\partial x} & \frac{\partial C_2}{\partial y} \\ \frac{\partial C_3}{\partial x} & \frac{\partial C_3}{\partial y} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \int_{I^2} \sqrt{|\frac{\partial C}{\partial x}|^2 + |\frac{\partial C}{\partial y}|^2 - \frac{\partial C}{\partial x} \frac{\partial C}{\partial y}}$$



On  $\mathbb{R}^3$

$$\mathcal{N}^0 \xrightarrow{d} \mathcal{N}^1 \xrightarrow{d} \mathcal{N}^2 \xrightarrow{d} \mathcal{N}^3$$





$$dF = W_{\text{grad} F} \quad dW_F^1 = W_{\text{curl} F}^2 \quad dW_G^2 = W_{\text{div} G}^3$$

Claim 1 IF  $N^1 \subset \mathbb{R}^3$ ,  $W_F^1 = (T \cdot F) ds$  on  $N^1$   
 oriented 1-manifold.

positive unit tangent to  $N$

the length form of  $N^1$

2 IF  $M^2 \subset \mathbb{R}^3$ ,  $W_G^2 = (G \cdot n) dA$  on  $M^2$   
 oriented 2-manifold

positive unit normal

Area Form

PF of 1 Eval both sides on  $T$ .

$$(T \cdot F) ds(T) = T \cdot F \cdot W_F^1(T)$$

$$= (F_1 dx_1 + F_2 dx_2 + F_3 dx_3) (T_1 e_1 + T_2 e_2 + T_3 e_3)$$


$$= F_1 T_1 + F_2 T_2 + F_3 T_3 = F \cdot T \quad \square$$

PF 2  $N^2$  IS  $W_G^2 = (G \cdot n) dA$  Enough to verify on  $(u, v)$ , a positive basis of  $T_p M$

$$W_G^2(u, v) = (G_1 dx_1 + dx_2 + dx_3 + \dots) \left( \begin{pmatrix} u_1 \\ u_2 \\ u_3 \end{pmatrix}, \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} \right) =$$

$$= G \cdot (u \times v) = G \cdot (|u \times v| \cdot n)$$

$$= (G \cdot n) |u \times v| = G \cdot n \cdot dA(u, v) \quad \square$$



$$\text{Area}_{xy}(u, v) = (u \times v) \cdot n \quad n = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

$u, v, w$  make an ON. basis.


$$\begin{aligned} A(\text{Proj}_{xy} C) &= (u \times v) \cdot n + (v \times w) \cdot n + (w \times u) \cdot n \\ &= (u + v + w) \cdot n \\ &= \text{length}. \end{aligned}$$

In  $\mathbb{R}^3$ :

1. on  $N^1 \subset \mathbb{R}^3$ ,  $W_F^1 = (T \cdot F) ds$

2. on  $M^2 \subset \mathbb{R}^3$ ,  $W_G^2 = (G \cdot n) dA$

$$\int_M dw = \int_{\partial M} w$$



Case 3  $M = M^3$   $W = W_G^2$

$$dW_F^1 = W_{\text{curl} F}^2$$

$$dw = W_{\text{div} G}^3 = (\text{div} G) (dx dy dz)$$

$$\int_M dw = \int_{\partial M} w$$


||

$$\int_M (\text{div} G) dV = \int_{\partial M} (G \cdot n) dA$$

"Gauss' thm"

$$\int \left( \frac{\partial G_1}{\partial x} + \frac{\partial G_2}{\partial y} + \frac{\partial G_3}{\partial z} \right) dV = \int_{\partial M} (G \cdot n) dA$$

The normal part of the flow



How much Flow is "created" within  $M$  = total outflow.

$$M = M^2 \quad W = W'_F \quad dW = W'^2_{\text{curl} F}$$


$$\int_M dW = \int_{\partial M} W$$

$$\int_M W'^2_{\text{curl} F} = \int_{\partial M} W'_F$$

$$\int_M (\text{curl} F) \cdot n \, dA = \int_{\partial M} (F \cdot T) \, ds \quad \text{"Stokes' theorem"}$$

$$\text{curl} F = \begin{pmatrix} \partial_2 F_3 - \partial_3 F_2 \\ \partial_3 F_1 - \partial_1 F_3 \\ \partial_1 F_2 - \partial_2 F_1 \end{pmatrix}$$

The rotation of a ball carried by the flow / whirling of the flow



$$\int_M (\text{curl} F) \cdot n \, dA = \int_{\partial M} (F \cdot T) \, ds$$

