

M.Sc. Math Workshop — Assignment #3

HUJI Spring 1998

Dror Bar-Natan

- (15) Prove: If $\lambda > 0$ is irrational and $\epsilon > 0$ then there exists 5 continuous functions $\phi_i : [0, 1] \rightarrow [0, 1]$ ($1 \leq i \leq 5$) so that for every continuous function $f : [0, 1] \times [0, 1] \rightarrow \mathbf{R}$ there exists a continuous function $g : [0, 1 + \lambda] \rightarrow \mathbf{R}$ so that

$$|f(x, y) - \sum_{i=1}^5 g(\phi_i(x) + \lambda\phi_i(y))| < \left(\frac{2}{3} + \epsilon\right) |f(x, y)|$$

for every $x, y \in [0, 1]$.

- (16) A *quantum probability space* is a pair (\mathcal{H}, v) where \mathcal{H} is a Hilbert space and $v \in \mathcal{H}$ is a unit vector. A *random variable* on \mathcal{H} is a self-adjoint operator $\mathcal{H} \rightarrow \mathcal{H}$. We say that $\langle v, A^n v \rangle$ is the expectation value of the n th power of the random variable A (if this quantity exists, namely if v is in the domain of definition of A^n). In particular, we set $E(A) = \langle v, Av \rangle$ to be the *expectation* of A , and $V(A) = \langle v, A^2 v \rangle - \langle v, Av \rangle^2$ to be the *variance* of A . Prove that if P and Q are random variables on some quantum probability space (\mathcal{H}, v) , and P and Q satisfy $[P, Q] = PQ - QP = iI$, then $V(P)V(Q) \geq \frac{1}{4}$. It is a good idea to start with the simplifying assumption $E(P) = E(Q) = 0$.
- (17) Prove that a finite group of affine transformations always has a fixed point.
- (18) Prove that the area of any planar section of a perfect tetrahedron is at most the area of a face of that tetrahedron.
- (19) Prove that any knot in \mathbf{R}^3 is the boundary of some double-sided (non-Möbius) surface embedded in \mathbf{R}^3 .
- (20) A rectangle R is tiled (presented as a disjoint union, not minding about 1-dimensional boundaries) with (possibly different) semi-integral rectangles — rectangles at least one of whose sides is of integral length. Prove that R itself is semi-integral.