



Dror Bar-Natan: Talks: GWU-1612:  
**On Elves and Invariants**



Follows Rozansky [Ro1, Ro2, Ro3] and Overbay [Ov], joint with van der Veen.

onef:=http://drorbn.net/GWU-1612/ 

Work in Progress! Fluid! Help Needed!

**Abstract.** Whether or not you like the formulas on this page, they describe the strongest truly computable knot invariant we know.

**Three steps to the computation of  $\rho_1$ :**

**1. Preparation.** Given  $K$ , results  $\langle \text{long word} \parallel \text{simple formulas} \rangle$ .

**2. Rewrite rules.** Make the word simpler and the formulas more complicated, until the word "elf" is reached.

**3. Readout.** The invariant  $\rho_1$  is read from the last formulas.

Knot  $K$

↓ preparation

$\langle \text{elf} \dots \text{elf} \parallel \omega_0; L_0; Q_0; P_0 \rangle$

↓ rewrite rules

$\langle \text{elf} \parallel \omega; -; -; P \rangle$

↓ readout

$\rho_1(K) = \rho_1(\omega, P)$

**Preparation.** Draw  $K$  using a 0-framed 0-rotation planar diagram  $D$  where all crossings are pointing up. Walk along  $D$  labeling features by  $1, \dots, m$  in order: over-passes, under-passes, and right-heading cups and caps (" $\pm$ -cuaps"). If  $x$  is a xing, let  $i_x$  and  $j_x$  be the labels on its over/under strands, and let  $s_x$  be 0 if it right-handed and  $-1$  otherwise. If  $c$  is a cuap, let  $i_c$  be its labels and  $s_c$  be its sign. Set

$$(L; Q; P) = \sum_{x: (i, j, s)} (-1)^s \left( l_j; t^s e_i f_j; (-1)^s e_i l_{(1+s)i-s} f_j + l_i l_j + \frac{t^{2s} e_i^2 f_j^2}{4} \right) + \sum_{c: (i, s)} (0; 0; s \cdot l_i).$$

This done, output  $\langle e_1 l_1 f_1 e_2 l_2 f_2 \dots e_m l_m f_m \parallel 1; L; Q; P \rangle$ .

**In formulas,**  $L$  is always  $\mathbb{Z}$ -linear in  $\{l_i\}$ ,  $Q$  is an  $R$ -linear combination of  $\{e_i f_j\}$  where  $R := \mathbb{Z}[t^{\pm 1}]$ , and  $P$  is an  $R$ -linear combination of  $\{1, l_i, l_i l_j, e_i f_j, e_i l_j f_k, e_i e_j f_k f_l\}$ .

**Rewrite Rules.** Manipulate  $\langle \text{word} \parallel \text{formulas} \rangle$  expressions using the rewrite rules below, until you come to the form  $\langle e_1 l_1 f_1 \parallel \omega; -; -; P \rangle$ . Output  $(\omega, P)$ .

**Rule 1, Deletions.** If a letter appears in word but not in formulas, you can delete it.

**Rule 2, Merges.** In word, you can replace adjacent  $v_i v_j$  with  $v_k$  (for  $v \in \{e, l, f\}$ ) while making the same changes in formulas (provided  $k$  creates no naming clashes). E.g.,


$$\langle \dots e_i e_j \dots \parallel Z \rangle \rightarrow \langle \dots e_k \dots \parallel Z_{|e_i e_j \rightarrow e_k} \rangle.$$

**Rule 3, le Sorts.** Provided  $k$  introduces no clashes, given  $\langle \dots l_j e_i \dots \parallel \omega; L; Q; P \rangle$ , decompose  $L = \lambda l_j + L'$ ,  $Q = \alpha e_i + Q'$ , write  $P = P(e_i, l_j)$  (with messy coefficients), set  $q = \epsilon^2 \beta e_k + \gamma l_k$ , and output


$$\langle \dots e_k l_k \dots \parallel \omega; L|_{l_j \rightarrow l_k}; t^q \alpha e_k + Q'; \epsilon^{-q} P(\partial_\alpha, \partial_\gamma) \epsilon^q |_{\beta \rightarrow \alpha/\omega, \gamma \rightarrow \lambda \log t} \rangle.$$

**Rule 4, fl Sorts.** Provided  $k$  introduces no clashes, given  $\langle \dots f_i l_j \dots \parallel \omega; L; Q; P \rangle$ , decompose  $L = \lambda l_j + L'$ ,  $Q = \alpha f_i + Q'$ , write  $P = P(f_i, l_j)$  (with messy coefficients), set  $q = \epsilon^2 \beta f_k + \gamma l_k$ , and output

$$\langle \dots l_k f_k \dots \parallel \omega; L|_{l_j \rightarrow l_k}; t^q \alpha f_k + Q'; \epsilon^{-q} P(\partial_\beta, \partial_\gamma) \epsilon^q |_{\beta \rightarrow \alpha/\omega, \gamma \rightarrow \lambda \log t} \rangle.$$



Happy Birthday, Scott!



The Knot Atlas

"God created the knots, all else in topology is the work of mortals."  
 Leopold Kronecker (modified) [www.katlas.org](http://www.katlas.org)

**Rule 5, fe Sorts.** Provided  $k$  introduces no clashes, given  $\langle \dots f_i e_j \dots \parallel \omega; L; Q; P \rangle$ , decompose  $Q = Q_{f_i e_j} + Q_{f_i} + Q_{e_j} + Q'$  write  $P = P(f_i, e_j)$  (with messy coefficients), set  $\mu = 1 + (t-1)\delta$  and  $q = ((1-t)\alpha\beta + \beta e_k + \alpha f_k + \delta e_k f_k)/\mu$ , and output

$$\left\langle \dots e_k f_k \dots \parallel \begin{matrix} \mu\omega; L; \mu\omega q + \mu Q' \\ \omega^4 \Lambda_k + \epsilon^{-q} P(\partial_\alpha, \partial_\beta) \epsilon^q \end{matrix} \right\rangle_{\substack{\alpha \rightarrow Q_{f_i e_j} / \omega, \beta \rightarrow Q_{f_i} / \omega, \\ \delta \rightarrow Q_{e_j} / \omega}}$$

where  $\Lambda_k$  is the  $\Delta\delta\gamma\alpha\zeta$ , "a principle of order and knowledge":

$$\Lambda_k = \frac{t+1}{4} \left( -\delta(\mu+1)(\beta^2 e_k^2 + \alpha^2 f_k^2) - \delta^3(3\mu+1)e_k^2 f_k^2 - 2(\beta e_k + \alpha f_k)(\alpha\beta + 2\delta\mu + \delta^2(2\mu+1)e_k f_k + 2\delta\mu^2 l_k) - 4(\alpha\beta + \delta\mu)(\delta(\mu+1)e_k f_k + \mu^2 l_k) - 4\delta^2 \mu^2 e_k f_k l_k + (t-1)(2(\alpha\beta + \delta\mu)^2 - \alpha^2 \beta^2) \right).$$

**elf merges,**  $m_k^{ij}$ , are defined as compositions


$$e_i l_i f_i e_j l_j f_j \xrightarrow{s_i^{f_i e_j}} e_i l_i e_x f_x l_j f_j \xrightarrow{s_i^{f_i e_x} \beta s_j^{f_i l_j}} e_i e_x l_x l_x f_x f_j \xrightarrow{l_j s_x \rightarrow k} e_k l_k f_k$$

**Readout.** Given  $\langle \text{elf} \parallel \omega; -; -; P \rangle$ , output

$$\rho_1(K) := \frac{(t\omega^3 - P|_{e, l, f=0})}{(t-1)^2 \omega^2}.$$

( $\omega$  is the Alexander polynomial,  $L$  and  $Q$  are not interesting.)

**Experimental Analysis ( $\omega\epsilon\beta/\text{Exp}$ ).** Log-log plot of computation time (sec) vs. crossing number, for all knots with up to 12 crossings (mean times) and for all torus knots with up to 48 crossings:



**Power.** On the 250 knots with at most 10 crossings, the pair  $(\omega, \rho_1)$  attains 250 distinct values, while (Khovanov, HOMFLY-PT) attains only 249 distinct values. To 11 crossings the numbers are (802, 788, 772) and to 12 they are (2978, 2883, 2786).

**Genus.** Up to 12 xings, always  $\rho_1$  is symmetric under  $t \leftrightarrow t^{-1}$ . With  $\rho_1^+$  denoting the positive-degree part of  $\rho_1$ , always  $\deg \rho_1^+ \leq 2g - 1$ , where  $g$  is the 3-genus of  $K$  (equality for 2530 knots). This gives a lower bound on  $g$  in terms of  $\rho_1$  (conjectural, but undoubtedly true). This bound is often weaker than the Alexander bound, yet for 10 of the 12-xing Alexander failures it does give the right answer.

**Why Works?** The Lie algebra  $\mathfrak{g}_1$  (below) is a "solvable approximation of  $\mathfrak{sl}_2$ ".

**Theorem.** The map (as defined below)

$$\langle w \parallel \omega; L; Q; P \rangle \mapsto \mathbb{C} \left( \omega^{-1} \epsilon^{L \log t + \omega^{-1} Q} (1 + \epsilon \omega^{-4} P) : w \right) \in \mathcal{U}(\mathfrak{g}_1)$$

is well defined modulo the sorting rules. It maps the initial preparation to a product of " $R$ -matrices" and "cuap values" satisfying the usual moves for Morse knots (R3, etc.). (And hence the result is a "quantum invariant", except computed very differently; no representation theory!).

Include Verification!

GWU-1612 Page 1

Totally rewrite, as In Talking Points on previous page.


**1-Smidgen  $sl_2$**  Let  $\mathfrak{g}_1$  be the 4-dimensional Lie algebra  $\mathfrak{g}_1 = \langle h, e', l, f \rangle$  over the ring  $R = \mathbb{Q}[\epsilon]/(\epsilon^2 = 0)$ , with  $h$  central and with  $[f, l] = f$ ,  $[e', l] = -e'$ , and  $[e', f] = h - 2\epsilon l$ . Over  $\mathbb{Q}$ ,  $\mathfrak{g}_1$  is a solvable approximation of  $sl_2$ :  $\mathfrak{g}_1 \supset \langle h, e', f, \epsilon h, \epsilon e', \epsilon l, \epsilon f \rangle \supset \langle h, \epsilon h, \epsilon e', \epsilon l, \epsilon f \rangle \supset 0$ . Pragmatics: declare  $\deg(h, e', l, f, \epsilon) = (1, 1, 0, 0, 1)$  and set  $t := e^h$  and  $e := (t-1)e'/h$ .

**How did it arise?**  $sl_2 = \mathfrak{b}^+ \oplus \mathfrak{b}^-/\mathfrak{b} := sl_2^+/ \mathfrak{b}$ , where  $\mathfrak{b}^+ = \langle l, f \rangle / [f, l] = f$  is a Lie bialgebra with  $\delta: \mathfrak{b}^+ \rightarrow \mathfrak{b}^+ \otimes \mathfrak{b}^+$  by  $\delta: (l, f) \mapsto (0, l \wedge f)$ . Going back,  $sl_2^+ = \mathcal{D}(\mathfrak{b}^+) = (\mathfrak{b}^+)^* \oplus \mathfrak{b}^+ = \langle h', e', l, f \rangle / \dots$ . **Idea.** Replace  $\delta \rightarrow \epsilon \delta$  over  $\mathbb{Q}[\epsilon]/(\epsilon^{k+1} = 0)$ . At  $k = 1$ , get  $[f, l] = f$ ,  $[f, h'] = -\epsilon f$ ,  $[l, e'] = e'$ ,  $[h', e'] = -\epsilon e'$ ,  $[h', l] = 0$ , and  $[e', f] = h' - \epsilon l$ . Now note that  $h' + \epsilon l$  is central, so switch to  $h := h' + \epsilon l$ . This is  $\mathfrak{g}_1$ .

**Ordering Symbols.**  $\odot$  (*poly* | *specs*) plants the variables of *poly* in  $\hat{\mathcal{S}}(\oplus_{\mathfrak{g}})$  along  $\hat{\mathcal{U}}(\mathfrak{g})$  according to *specs*. E.g.,  $\odot(e_1 e^3 f_1^2 f_2^3 | f_3 l_1 e_1 e_3 l_2) = f^3 l^2 e e^3 l \in \hat{\mathcal{U}}(\mathfrak{g})$ .

This enables the description of elements of  $\hat{\mathcal{U}}(\mathfrak{g})$  using commutative polynomials / power series. In  $\mathfrak{g}_1$ , no need to specify  $h/t$ .

**Algebras and Invariants.** Given any unital algebra  $A$  (even better if  $A$  is Hopf; typically,  $A \sim \hat{\mathcal{U}}(\mathfrak{g})$ ), appropriate orange  $R \in A \otimes A$ , and appropriate cuaps  $\in A$ , get an  $A^{\otimes S}$ -valued invariant of pure  $S$ -component tangles:



**What we didn't say** (more, including videos, in  $\omega\epsilon\beta$ /Talks).

- $\rho_1$  is "line" in the coloured Jones polynomial; related to Melvin-Morton-Rozansky.
- $\rho_1$  extends to "rotational virtual tangles" and is a projection of the universal finite type invariant of such.
- $\rho_1$  seems to have a better chance than anything else we know to detect a counterexample to slice=ribbon.
- $\rho_1$  leads to many questions and a very long to-do list. Years of work, many papers ahead. Have fun!

**Demo Programs.**  $\omega\epsilon\beta$ /Demo

```
CF[e_] := Module[{vars = Union@Cases[e, e_ | 1_ | f_, {}],
  If[vars == {}, Factor[e],
  Total[CoefficientRules[e, vars] /
  (p_ -> c_) => Factor[c] Times @@ (vars^p) ]];];
CF[e_E] := CF[e];
```

```
E[i_, j_, s_] := E[1, (-1)^s 1_], (-1)^s e_i f_j,
t^s e_i 1_{(1+s) i-s j} f_j + (-1)^s 1_i 1_j + (-t^2)^s e_i^2 f_j^2 / 4];
E[i_, s_] := E[1, 0, 0, s 1_i];
E /: E[1, L1_, Q1_, P1_] E[1, L2_, Q2_, P2_] :=
E[1, L1 + L2, Q1 + Q2, P1 + P2];
```

**Preparation**

**z1 = (E[1, 11, 0] E[4, 2, -1] E[15, 5, 0] Preparing the Trefoil**  
**E[6, 8, -1] E[9, 16, 0] E[12, 14, -1] E[3, -1] E[7, +1]**  
**E[10, -1] E[13, +1])**

$$E[1, -1_2 + 1_5 - 1_8 + 1_{11} - 1_{14} + 1_{16},$$

$$- \frac{e_4 f_2}{t} + e_{15} f_5 - \frac{e_6 f_8}{t} + e_1 f_{11} - \frac{e_{12} f_{14}}{t} + e_9 f_{16},$$

$$- \frac{e_4^2 f_2^2}{4 t^2} + \frac{1}{4} e_{15}^2 f_5^2 - \frac{e_6^2 f_8^2}{4 t^2} + \frac{1}{4} e_1^2 f_{11}^2 - \frac{e_{12}^2 f_{14}^2}{4 t^2} + \frac{1}{4} e_9^2 f_{16}^2 + e_1 f_{11} 1_1 +$$

$$\frac{e_4 f_2 1_2}{t} - 1_3 - 1_2 1_4 + 1_7 + \frac{e_6 f_8 1_8}{t} - 1_6 1_8 + e_9 f_{16} 1_9 - 1_{18} +$$

$$1_1 1_{11} + 1_{13} + \frac{e_{12} f_{14} 1_{14}}{t} - 1_{12} 1_{14} + e_{15} f_5 1_{15} + 1_5 1_{15} + 1_9 1_{16}]$$

**DP<sub>x->0, y->0, z->0</sub>[P\_][f\_]** := **Differential Polynomials**

**Total[CoefficientRules[P, {x, y}]] /.** (Implementing  $P(\partial_x, \partial_y)(f)$ )  
 $\{m_-, n_-\} \rightarrow c_-\} \Rightarrow CD[f, \{a, m\}, \{b, n\}]$

```
S1_j_[{x:e|f}_i_>h_>R_][E[omega_-, L_-, Q_-, P_]] := le and fl Sorts
With[{lambda = theta_1_j L_-, alpha = theta_x_i Q_-, q = e^gamma beta x_h + gamma 1_h}, CF[
E[omega_-, L_ / . 1_j -> 1_h, t^lambda alpha x_h + (Q / . x_i -> theta),
e^-q DP_{1_j->0, x_i->0}[P][e^q] / . {beta -> alpha / omega, gamma -> lambda Log[t]} ]];];
```

$$\Lambda[h_-] := ((t-1) (2(\alpha\beta + \delta\mu)^2 - \alpha^2\beta^2) - 4e_h 1_h f_h \delta^2 \mu^2 -$$

$$\delta(1+\mu) (f_h^2 \alpha^2 + e_h^2 \beta^2) - e_h^2 f_h^2 \delta^3 (1+3\mu) -$$

$$2(\alpha\beta + 2\delta\mu + e_h f_h \delta^2 (1+2\mu) + 2 1_h \delta \mu^2) (f_h \alpha + e_h \beta) -$$

$$4(1_h \mu^2 + e_h f_h \delta (1+\mu)) (\alpha\beta + \delta\mu)) (1+t) / 4;$$

**S<sub>f\_i e\_j -> h</sub>[E[omega\_-, L\_-, Q\_-, P\_]]** := **fe Sorts**

```
With[{q = ((1-t) alpha beta + beta e_h + alpha f_h + delta e_h f_h) / mu}, CF[
E[mu omega_-, L_-, mu omega q + mu (Q / . f_i | e_j -> theta),
mu^4 e^-q DP_{f_i->0, e_j->0}[P][e^q] + omega^A Lambda[h_-]] / . mu -> 1 + (t-1) delta / .
{alpha -> omega^-1 (theta_i Q / . e_j -> theta), beta -> omega^-1 (theta_j Q / . f_i -> theta),
delta -> omega^-1 theta_{f_i, e_j Q}}];];
```

**m\_{i->j, h}**[Z\_E] := **Module**[{x, z}, **Elf Merges**

```
CF[Z // S_{f_i e_j -> h} // S_{1_i e_h -> x} // S_{f_h 1_j -> x} / . z_{-1|j}|x -> z_h]]
```

**(Do[z1 = z1 // m\_{1, k-1}, {k, 2, 16}]; z1)** **Rewriting the Trefoil**

$$E\left[\frac{1-t-t^2}{t}, \theta, \theta, \frac{(1-t)t}{t^3} \frac{(1-t+t^2)^2}{t^3} \frac{(1-t-2t^2)}{t^4} -$$

$$\frac{2(1-t)(1-t+t^2)^3 e_1 f_1}{t^4} - \frac{2(-1-t)(1-t)(1-t+t^2)^3 1_1}{t^4}\right]$$

**rho\_1[E[omega\_-, \_ , \_ , P\_]]** := **CF**  $\left[\frac{t}{(t-1)^2 \omega^2} \left(\frac{P / . e_- | 1_- | f_- \rightarrow \theta}{t} - t \omega^2 (\theta_t \omega)\right)\right]$

**rho\_1[z1] // Expand** **rho\_1(31)**

$\frac{1}{t} + t$

**References.**

[Ov] A. Overbay, *Perturbative Expansion of the Colored Jones Polynomial*, University of North Carolina PhD thesis,  $\omega\epsilon\beta$ /Ov.  
[Ro1] L. Rozansky, *A contribution of the trivial flat connection to the Jones polynomial and Witten's invariant of 3d manifolds*, I. Comm. Math. Phys. 175-2 (1996) 275-296, arXiv:hep-th/9401061.  
[Ro2] L. Rozansky, *The Universal R-Matrix, Burau Representation and the Melvin-Morton Expansion of the Colored Jones Polynomial*, Adv. Math. 134-1 (1998) 1-31, arXiv:q-alg/9604005.  
[Ro3] L. Rozansky, *A Universal U(1)-RCC Invariant of Links and Rationality Conjecture*, arXiv:math/0201139.

better naming!

diagram	$n_i^+$	Alexander's $\omega^-$	Today's / Rozansky's $\rho_1^+$	unknotting number / amphicheiral	genus / ribbon	diagram	$n_i^+$	Alexander's $\omega^-$	Today's / Rozansky's $\rho_1^+$	unknotting number / amphicheiral	genus / ribbon
	$0_1^+$	1		0 / ✓	0 / ✓		$3_1^+$	$t-1$		1 / ✗	1 / ✗
	$4_1^+$	$3-t$		1 / ✗	1 / ✓		$5_1^+$	$t^2-t+1$		2 / ✗	2 / ✗
	$0$						$2_1^+$	$t^3+3t$			2 / ✗