




# Algebraic Knot Theory




**Abstract.** This will be a very “light” talk: I will explain why about 13 years ago, in order to have a say on some problems in knot theory, I’ve set out to find tangle invariants with some nice compositional properties. I will not mention that in recent joint work with Roland van der Veen we’ve found such invariants, and we are now struggling to make use of them.

## (v-)Tangles.



**Genus.** Every knot is the boundary of an orientable “Seifert Surface” (ωεβ/SS), and the least of their genera is the “genus” of the knot.

**Claim.** The knots of genus ≤ 2 are precisely the images of 4-component tangles via




**A Bit about Ribbon Knots.** A “ribbon knot” is a knot that can be presented as the boundary of a disk that has “ribbon singularities”, but no “clasp singularities”. A “slice knot” is a knot in  $S^3 = \partial B^4$  which is the boundary of a non-singular disk in  $B^4$ . Every ribbon knot is clearly slice, yet,

**Conjecture.** Some slice knots are not ribbon.

**Fox-Milnor.** The Alexander polynomial of a ribbon knot is always of the form  $A(t) = f(t)f(1/t)$ . (also for slice)

**Theorem.**  $K$  is ribbon iff it is  $\kappa T$  for a tangle  $T$  for which  $\tau T$  is the untangle  $U$ .



**The Gold Standard** is set by the “T-calculus” Alexander formulas [BNS, BN]. An  $S$ -component tangle  $T$  has  $\Gamma(T) \in R_S \times M_{S \times S}(R_S) = \left\{ \begin{array}{c|c} \omega & S \\ \hline S & A \end{array} \right\}$  with  $R_S := \mathbb{Z}\langle\langle T_a : a \in S \rangle\rangle$ :

$$\begin{aligned} (a \nearrow b, b \nearrow a) &\rightarrow \begin{array}{c|c|c} 1 & a & b \\ \hline a & 1 & 1 - T_a^{-1} \\ b & 0 & T_a^{-1} \end{array} & T_1 \sqcup T_2 &\rightarrow \begin{array}{c|c|c|c} \omega_1 \omega_2 & S_1 & S_2 & \\ \hline S_1 & A_1 & 0 & \\ S_2 & 0 & A_2 & \end{array} \\ \begin{array}{c|c|c} \omega & a & b & S \\ \hline a & \alpha & \beta & \theta \\ b & \gamma & \delta & \epsilon \\ S & \phi & \psi & \Xi \end{array} & \xrightarrow{m_c^{ab}} & \begin{array}{c|c|c} (1-\beta)\omega & c & S \\ \hline c & \gamma + \frac{\alpha\delta}{1-\beta} & \epsilon + \frac{\delta\theta}{1-\beta} \\ S & \phi + \frac{\alpha\psi}{1-\beta} & \Xi + \frac{\psi\theta}{1-\beta} \end{array} \end{aligned}$$

For long knots,  $\omega$  is Alexander, and that’s the fastest Alexander algorithm I know! Dunfield: 1000-crossing fast.

## Strand Doubling and Reversal.

$$\begin{array}{c|c|c} \omega & a & S \\ \hline a & \alpha & \theta \\ S & \phi & \Xi \end{array} \xrightarrow{\begin{array}{l} q\Delta_{bc}^a \\ \mu T_a^{-1} \\ \nu T_a^{-1} \\ T_a^{-1} T_b^{-1} T_c^{-1} \end{array}} \begin{array}{c|c|c|c} \omega & b & c & S \\ \hline b & (\sigma_a - \alpha T_a - \nu T_c)/\mu & (T_b - 1)T_c \nu/\mu & (T_b - 1)T_c \theta/\mu \\ c & (T_c - 1)\nu/\mu & (\alpha - \sigma_a T_a - \nu T_c)/\mu & (T_c - 1)\theta/\mu \\ S & \phi & \phi & \Xi \end{array}$$

**Vo’s Thesis [Vo].** A proof of the Fox-Milnor theorem for ribbon knots using this technology (and more).

**Implementation key idea:** ωεβ/AlexDemo

```
Γ := Γ[ω1, λ1, ω2, λ2] := Γ[ω1*ω2, λ1+λ2];
Module[α, β, γ, δ, θ, ε, φ, ψ, Ξ, μ];
Collect[Γ[ω, λ], h, Collect[#, t, Factor] &];
Format[Γ[ω, λ]] := Module[{S, M},
M = Outer[Factor][θ, ω, λ] & S, S, S];
M = Prepend[M, t & /@ S] // Transpose;
M = Prepend[M, Prepend[h & /@ S, ω]];
M // MatrixForm;
```

**Meta-Associativity**  $\xi = \Gamma[\omega, \{t_1, t_2, t_3, t_s\}] \cdot \begin{pmatrix} \alpha_{11} & \alpha_{12} & \alpha_{13} & \theta_1 \\ \alpha_{21} & \alpha_{22} & \alpha_{23} & \theta_2 \\ \alpha_{31} & \alpha_{32} & \alpha_{33} & \theta_3 \\ \phi_1 & \phi_2 & \phi_3 & \Xi \end{pmatrix} \cdot \{h_1, h_2, h_3, h_s\}$

$$(\xi // m_{12 \rightarrow 1} // m_{13 \rightarrow 1}) = (\xi // m_{23 \rightarrow 2} // m_{12 \rightarrow 1})$$

**True R3** ... divide and conquer!

$$\begin{pmatrix} 1 & h_1 & h_2 & h_3 \\ t_1 & \frac{T_3}{T_2} & 0 & 0 \\ t_2 & \frac{-1+T_2}{T_2} & \frac{1}{T_3} & 0 \\ t_3 & \frac{-1+T_3}{T_2} & \frac{-1+T_3}{T_3} & 1 \end{pmatrix}, \begin{pmatrix} 1 & h_1 & h_2 & h_3 \\ t_1 & \frac{T_3}{T_2} & 0 & 0 \\ t_2 & \frac{-1+T_2}{T_2} & \frac{1}{T_3} & 0 \\ t_3 & \frac{-1+T_3}{T_2} & \frac{-1+T_3}{T_3} & 1 \end{pmatrix}$$

**Do** [z = z // m<sub>1k-1</sub>, {k, 2, 16}];

$$\begin{pmatrix} 11 - \frac{1}{T_1} + \frac{4}{T_1^2} - \frac{8}{T_1} - 8 T_1 + 4 T_1^2 - T_1^3 & & & \\ & t_1 & & \\ & & & 1 \end{pmatrix}$$

**Fact.**  $\Gamma$  is better viewed as an invariant of a certain class of 2D knotted objects in  $\mathbb{R}^4$  [BND, BN].


**Fact.**  $\Gamma$  is the “0-loop” part of an invariant that generalizes to “n-loops” (1D tangles only, see ωεβ/talks and future publications with van der Veen).

**Speculation.** Stepping stones to categorification?

[BN] D. Bar-Natan, *Balloons and Hoops and their Universal Finite Type Invariant, BF Theory, and an Ultimate Alexander Invariant*, ωεβ/KBH, arXiv:1308.1721.  
[BND] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects I: w-Knots and the Alexander Polynomial*, Alg. and Geom. Top. **16-2** (2016) 1063–1133, arXiv:1405.1956, ωεβ/WKO1.  
[BNS] D. Bar-Natan and S. Selmani, *Meta-Monoids, Meta-Bicrossed Products, and the Alexander Polynomial*, J. of Knot Theory and its Ramifications **22-10** (2013), arXiv:1302.5689.  
[GST] R. E. Gompf, M. Scharlemann, and A. Thompson, *Fibered Knots and Potential Counterexamples to the Property 2R and Slice-Ribbon Conjectures*, Geom. and Top. **14** (2010) 2305–2347, arXiv:1103.1601.  
[Vo] H. Vo, *Alexander Invariants of Tangles via Expansions*, University of Toronto Ph.D. thesis, ωεβ/Vo.

“God created the knots, all else in topology is the work of mortals.”  
Leopold Kronecker (modified)

# Proof of the Tangle Characterization of Ribbon Knots




**Theorem.** A knot  $K$  is ribbon iff there exists a tangle  $T$  whose  $\tau$  closure is the untangle and whose  $\kappa$  closure is  $K$ .

**Proof.** The backward  $\Leftarrow$  implication is easy:




For the forward implication, follow the following 5 steps:




Step 1: In-situ cosmetics.

At end:  $D$  is a tree of chord-and-arc polygons.




Step 2: Near-situ cosmetics.

At end:  $D$  is tree-band-sum of  $n$  unknotted disks.



Step 3: Slides.


At end:  $D$  is a linear-band-sum of  $n$  unknotted disks.



Step 4: Exposure!

The green domain is contractible - so it can be shrunk, moved at will (with the blue membrane following along), and expanded back again.

At end:  $D$  has  $(n-1)$  exposed bridges which when turned, make  $D$  a union of  $n$  unknotted disks.



Step 5: Pulling bottom handles avoiding the obstacles.

At end: Theorem is proven.

