

Pensieve header: A talk and a program about Archibald- (\mathcal{A} -) and Γ -calculus and the Halacheva map between them; the \mathcal{A} part. Continues pensieve://2021-03/

\mathcal{A} -Calculus

tex

```
\begin{frame}
{\LARGE 3. An Implementation of  $\mathcal{A}$ }
If I didn't implement I wouldn't believe myself.
\vskip 2mm
Written in Mathematica~\cite{Wolfram:Mathematica}, available as the notebook {\tt Alpha.nb} at
\url{http://drorbn.net/mo21/ap}. Code lines are highlighted in grey, demo lines are plain.
We start with an implementation of elements ( $\mathcal{W}$ ) of exterior algebras, and of the wedge
product ( $\mathcal{WP}$ ):
\vskip 2mm
```

pdf

```
In[ ]:= WP[Wedge[u___], Wedge[v___]] := Signature[{u, v}] * Wedge @@ Sort[{u, v}];
WP[0, _] = WP[_ , 0] = 0;
WP[A_, B_] :=
Expand[Distribute[A ** B] /. (a_. * u_Wedge) ** (b_. * v_Wedge) -> a b WP[u, v]];
```

pdf

```
In[ ]:= WP[Wedge[] + Wedge[a] - 2 b ^ a, Wedge[] - 3 Wedge[b] + 7 c ^ d]
```

Out[]:=

pdf

```
Wedge[] + Wedge[a] - 3 Wedge[b] - a ^ b + 7 c ^ d + 7 a ^ c ^ d + 14 a ^ b ^ c ^ d
```

tex

```
\end{frame}
\begin{frame}\null
We then define the exponentiation map in exterior algebras ( $\mathcal{WExp}$ ) by summing the series and
stopping the sum once the current term ( $\mathcal{t}$ ) vanishes:
```

pdf

```
In[ ]:= WExp[A_] := Module[{s = Wedge[], t = Wedge[], k = 0},
While[t != 0, s += (t = Expand[WP[t, A] / (++k)])]; s]
```

pdf

```
In[ ]:= WExp[a ^ b + c ^ d + e ^ f]
```

Out[]:=

pdf

```
Wedge[] + a ^ b + c ^ d + e ^ f + a ^ b ^ c ^ d + a ^ b ^ e ^ f + c ^ d ^ e ^ f + a ^ b ^ c ^ d ^ e ^ f
```

tex

```
\end{frame}
\begin{frame}\null
Contractions!
```

pdf

```
In[ ]:= c_{x,y}[w_Wedge] := Module[{i, j},
  {i} = FirstPosition[w, x, {0}]; {j} = FirstPosition[w, y, {0}];
  {
    w (i == 0) ^ (j == 0)
    (-1)^{i+j+If[i>j,0,1]} Delete[w, {{i}, {j}}] (i > 0) ^ (j > 0)
  };
  c_{x,y}[E_] := E /. w_Wedge -> c_{x,y}[w]
```

pdf

```
In[ ]:= WExp[a ^ b + 2 c ^ d]
Cd,c@WExp[a ^ b + 2 c ^ d]
```

Out[]= pdf

Wedge[] + a ^ b + 2 c ^ d + 2 a ^ b ^ c ^ d

Out[]= pdf

-Wedge[] - a ^ b

tex


\end{frame}

tex

\begin{frame}

\parpic[r]{\def\X{\mathcal{X}_{\{ijkl\}[S,T]}\input{figs/Xp.pdf_t}}

$\mathcal{X}_{\{ijkl\}[S,T]}$ is also a container for the values of the \mathcal{X} -invariant of a tangle. In it, $\{i, j, k, l\}$ are the labels of the input strands, $\{s, t\}$ are the labels of the output strands, $\{c\}$ is an assignment of colours (namely, variables) to all the ends $\{x_i\}_i \in \text{in}(\text{is}) \sqcup \{x_j\}_j \in \text{in}(\text{os})$, and $\{w\}$ is the "payload": an element of $\Lambda(\text{in}(\text{is}) \sqcup \text{in}(\text{os}))$.



pdf

```
In[ ]:= A[X_{i,j,k,l}[S_, T_]] := A[{L, i}, {j, k}, <| E_i -> S, x_j -> T, x_k -> S, E_l -> T |>,
  Expand[T^{-1/2} WExp[Expand[{E_l, E_i} . (1 1 - T; 0 T) . {x_j, x_k} /. E_a x_b -> E_a ^ x_b]]];
```

pdf

In[*]:= $\mathcal{A}[X_{1,2,3,4}[u, v]]$

Out[*]=
pdf

$$\mathcal{A}\left[\{4, 1\}, \{2, 3\}, \langle \xi_1 \rightarrow u, x_2 \rightarrow v, x_3 \rightarrow u, \xi_4 \rightarrow v \rangle, \frac{\text{Wedge}[] - \frac{x_2 \wedge \xi_4}{\sqrt{v}} - \sqrt{v} x_3 \wedge \xi_1 - \frac{x_3 \wedge \xi_4}{\sqrt{v}} + \sqrt{v} x_3 \wedge \xi_4 + \sqrt{v} x_2 \wedge x_3 \wedge \xi_1 \wedge \xi_4}{\sqrt{v}}\right]$$

pdf

In[*]:= $\mathcal{A}[X_{i,j,k,L}] := \mathcal{A}[X_{i,j,k,L}[\tau_i, \tau_L]]$

tex

`\end{frame}`

tex


`\begin{frame}\null`

The negative crossing and the “point”:

`\[\def\Xbar{\bar{X}}_{ijkl}[S,T] \def\P{P}_{ij}[T] \input{figs/XmP.pdf_t}`

`\input{figs/XmP.pdf_t}`

`\]`



pdf

In[*]:= $\mathcal{A}[\bar{X}_{i,j,k,L}[S_-, T_-]] := \mathcal{A}[\{i, j\}, \{k, l\}, \langle \xi_i \rightarrow S, \xi_j \rightarrow T, x_k \rightarrow S, x_l \rightarrow T \rangle, \text{Expand}\left[T^{1/2} \text{WExp}\left[\text{Expand}\left[\{\xi_i, \xi_j\} \cdot \begin{pmatrix} T^{-1} & 0 \\ 1 - T^{-1} & 1 \end{pmatrix} \cdot \{x_k, x_l\}\right] / \cdot \xi_a x_b \Rightarrow \xi_a \wedge x_b\right]\right]$
 $\mathcal{A}[\bar{X}_{i,j,k,L}] := \mathcal{A}[\bar{X}_{i,j,k,L}[\tau_i, \tau_j]]$

pdf

In[*]:= $\mathcal{A}[P_{i,j}[T_-]] := \mathcal{A}[\{i\}, \{j\}, \langle \xi_i \rightarrow T, x_j \rightarrow T \rangle, \text{WExp}[\xi_i \wedge x_j]]$
 $\mathcal{A}[P_{i,j}] := \mathcal{A}[P_{i,j}[\tau_i]]$

tex

`\end{frame}`

`\begin{frame}\null`

The linear structure on \mathcal{A} 's:

pdf

In[*]:= $\mathcal{A} /: \alpha \times \mathcal{A}[is_-, os_-, cs_-, w_-] := \mathcal{A}[is, os, cs, \text{Expand}[\alpha w]]$
 $\mathcal{A} /: \mathcal{A}[is1_-, os1_-, cs1_-, w1_-] + \mathcal{A}[is2_-, os2_-, cs2_-, w2_-] /;$
 $(\text{Sort}@is1 == \text{Sort}@is2) \wedge (\text{Sort}@os1 == \text{Sort}@os2) \wedge$
 $(\text{Sort}@Normal@cs1 == \text{Sort}@Normal@cs2) := \mathcal{A}[is1, os1, cs1, w1 + w2]$

tex

Deciding if two \mathcal{A} 's are equal:

pdf

```
In[*]:=  $\mathcal{A} /: \mathcal{A}[is1_, os1_, \_, w1_] \equiv \mathcal{A}[is2_, os2_, \_, w2_] :=$   

TrueQ[(Sort@is1 === Sort@is2) \wedge (Sort@os1 === Sort@os2) \wedge PowerExpand[w1 == w2]]
```

tex

```
\end{frame}  

\begin{frame}\null  

\vskip -12mm  

\null\hfill\smash{\imagetop{\input{figs/R2Left.pdf_t}}}  

\newline The union operation on  $\mathcal{A}$ 's (implemented as ``multiplication'')
```

pdf

```
In[*]:=  $\mathcal{A} /: \mathcal{A}[is1_, os1_, cs1_, w1_] \mathcal{A}[is2_, os2_, cs2_, w2_] :=$   

 $\mathcal{A}[is1 \cup is2, os1 \cup os2, Join[cs1, cs2], WP[w1, w2]]$ 
```

pdf

```
In[*]:= Short[ $\mathcal{A}[X_{2,4,3,1}[S, T]] \mathcal{A}[\bar{X}_{3,4,6,5}], 5]$ 
```

Out[*]//Short=

pdf

$$\mathcal{A}[\{1, 2, 3, 4\}, \{3, 4, 5, 6\}, \langle \xi_2 \rightarrow S, x_4 \rightarrow T, x_3 \rightarrow S, \xi_1 \rightarrow T, \xi_3 \rightarrow \tau_3, \xi_4 \rightarrow \tau_4, x_6 \rightarrow \tau_3, x_5 \rightarrow \tau_4 \rangle,$$

$$\frac{\sqrt{\tau_4} \text{Wedge}[]}{\sqrt{T}} - \frac{\sqrt{\tau_4} x_3 \wedge \xi_1}{\sqrt{T}} + \sqrt{T} \sqrt{\tau_4} x_3 \wedge \xi_1 - \sqrt{T} \sqrt{\tau_4} x_3 \wedge \xi_2 -$$

$$\frac{\sqrt{\tau_4} x_4 \wedge \xi_1}{\sqrt{T}} - \frac{\sqrt{\tau_4} x_5 \wedge \xi_4}{\sqrt{T}} - \frac{x_6 \wedge \xi_3}{\sqrt{T} \sqrt{\tau_4}} + \ll 40 \gg + \frac{\sqrt{T} x_3 \wedge x_5 \wedge x_6 \wedge \xi_1 \wedge \xi_3 \wedge \xi_4}{\sqrt{\tau_4}} -$$

$$\left. \frac{\sqrt{T} x_3 \wedge x_5 \wedge x_6 \wedge \xi_2 \wedge \xi_3 \wedge \xi_4}{\sqrt{\tau_4}} - \frac{x_4 \wedge x_5 \wedge x_6 \wedge \xi_1 \wedge \xi_3 \wedge \xi_4}{\sqrt{T} \sqrt{\tau_4}} + \frac{\sqrt{T} x_3 \wedge x_4 \wedge x_5 \wedge x_6 \wedge \xi_1 \wedge \xi_2 \wedge \xi_3 \wedge \xi_4}{\sqrt{\tau_4}} \right]$$

tex

```
\end{frame}  

\begin{frame}\null  

Contractions of  $\mathcal{A}$ -objects:
```

pdf

```
In[*]:=  $c_{h,t} @ \mathcal{A}[is_, os_, cs_, w_] := \mathcal{A}$   

DeleteCases[is, t], DeleteCases[os, h], KeyDrop[cs, {x_h, \xi_t}],  $c_{x_h, \xi_t}[w]$   

] /. If[MatchQ[cs[\xi_t], \tau_], cs[\xi_t] \rightarrow cs[x_h], cs[x_h] \rightarrow cs[\xi_t]];
```

pdf

In[*]:= $c_{4,4}[\mathcal{A}[X_{2,4,3,1}[S, T]] \mathcal{A}[\bar{X}_{3,4,6,5}]]$

Out[*]=

pdf

$$\mathcal{A}[\{1, 2, 3\}, \{3, 5, 6\}, \langle \xi_2 \rightarrow S, x_3 \rightarrow S, \xi_1 \rightarrow T, \xi_3 \rightarrow T, x_6 \rightarrow T, x_5 \rightarrow T \rangle, \\ \text{Wedge}[] - x_3 \wedge \xi_1 + T x_3 \wedge \xi_1 - T x_3 \wedge \xi_2 - x_5 \wedge \xi_1 - x_6 \wedge \xi_1 + \frac{x_6 \wedge \xi_1}{T} - \frac{x_6 \wedge \xi_3}{T} + \\ T x_3 \wedge x_5 \wedge \xi_1 \wedge \xi_2 - x_3 \wedge x_6 \wedge \xi_1 \wedge \xi_2 + T x_3 \wedge x_6 \wedge \xi_1 \wedge \xi_2 + x_3 \wedge x_6 \wedge \xi_1 \wedge \xi_3 - \\ \frac{x_3 \wedge x_6 \wedge \xi_1 \wedge \xi_3}{T} - x_3 \wedge x_6 \wedge \xi_2 \wedge \xi_3 - \frac{x_5 \wedge x_6 \wedge \xi_1 \wedge \xi_3}{T} - x_3 \wedge x_5 \wedge x_6 \wedge \xi_1 \wedge \xi_2 \wedge \xi_3]$$

tex

\end{frame}

\begin{frame}\null

Automatic and intelligent multiple contractions:

pdf

```
In[*]:= c@A[is_, os_, cs_, w_] := Fold[c_{#2, #2} [#1] &, A[is, os, cs, w], is ∩ os]
A[{A_ A_}] := c[A];
A[{A1_ A_, As_ A_}] := Module[{A2},
  A2 = First@MaximalBy[{As}, Length[A1[[1]] ∩ #[[2]]] + Length[A1[[2]] ∩ #[[1]]] &];
  A[Join[{c[A1 A2]}, DeleteCases[{As}, A2]]];
A[os_List] := A[A/@os]
```

pdf

In[*]:= $c[\mathcal{A}[X_{2,4,3,1}[S, T]] \mathcal{A}[\bar{X}_{3,4,6,5}]]$

Out[*]=

pdf

$$\mathcal{A}[\{1, 2\}, \{5, 6\}, \langle \xi_2 \rightarrow S, \xi_1 \rightarrow T, x_6 \rightarrow S, x_5 \rightarrow T \rangle, \text{Wedge}[] - x_5 \wedge \xi_1 - x_6 \wedge \xi_2 - x_5 \wedge x_6 \wedge \xi_1 \wedge \xi_2]$$

pdf

In[*]:= $\mathcal{A}@\{\mathcal{A}[X_{2,4,3,1}[S, T]], \mathcal{A}[\bar{X}_{3,4,6,5}]\}$

Out[*]=

pdf

$$\mathcal{A}[\{1, 2\}, \{5, 6\}, \langle \xi_2 \rightarrow S, \xi_1 \rightarrow T, x_6 \rightarrow S, x_5 \rightarrow T \rangle, \text{Wedge}[] - x_5 \wedge \xi_1 - x_6 \wedge \xi_2 - x_5 \wedge x_6 \wedge \xi_1 \wedge \xi_2]$$

tex

\end{frame}

Skein Relations



tex

\begin{frame}\{LARGE 4. Skein relations and evaluations for \mathcal{A} \}

`\[\input{figs/SimpleTangle.pdf_t} \]`

pdf

`In[*]:=` $\mathcal{A} @ \{ \bar{X}_{4,1,6,3} [v, u], \bar{X}_{3,2,5,4} \}$

`Out[*]=`
pdf

$$\mathcal{A} [\{ 1, 2 \}, \{ 5, 6 \}, \langle | \xi_2 \rightarrow v, x_5 \rightarrow u, \xi_1 \rightarrow u, x_6 \rightarrow v | \rangle ,$$

$$\sqrt{u} \sqrt{v} \text{Wedge} [] - \frac{\sqrt{u} x_5 \wedge \xi_1}{\sqrt{v}} + \frac{\sqrt{u} x_5 \wedge \xi_2}{\sqrt{v}} - \sqrt{u} \sqrt{v} x_5 \wedge \xi_2 + \frac{\sqrt{v} x_6 \wedge \xi_1}{\sqrt{u}} - \sqrt{u} \sqrt{v} x_6 \wedge \xi_1 -$$

$$\frac{\sqrt{v} x_6 \wedge \xi_2}{\sqrt{u}} - \frac{\sqrt{u} x_5 \wedge x_6 \wedge \xi_1 \wedge \xi_2}{\sqrt{v}} - \frac{\sqrt{v} x_5 \wedge x_6 \wedge \xi_1 \wedge \xi_2}{\sqrt{u}} + \sqrt{u} \sqrt{v} x_5 \wedge x_6 \wedge \xi_1 \wedge \xi_2]$$


Reidemeister 2

tex

`\end{frame}`
`\begin{frame}{\large Reidemeister 2}`

tex

`\[\input{figs/R2.pdf_t} \]`



pdf

`In[*]:=` $\mathcal{A} @ \{ X_{2,4,3,1} [S, T], \bar{X}_{3,4,6,5} \} \equiv \mathcal{A} @ \{ P_{1,5} [T], P_{2,6} [S] \}$

`Out[*]=`
pdf

True

pdf

`In[*]:=` $\mathcal{A} @ \{ \bar{X}_{3,1,2,4} [S, T], X_{6,5,3,4} \} \equiv \mathcal{A} @ \{ P_{1,5} [T], P_{6,2} [S] \}$

`Out[*]=`
pdf

True


Reidemeister 3

tex

`\end{frame}`
`\begin{frame}{\large Reidemeister 3}`

tex

`\[\input{figs/R3.pdf_t} \]`



pdf

In[*]:= $\mathcal{A}@\{X_{2,5,4,1}[T_2, T_1], X_{3,7,6,5}[T_3, T_1], X_{6,9,8,4}\} \equiv \mathcal{A}@\{X_{3,5,4,2}[T_3, T_2], X_{4,6,8,1}[T_3, T_1], X_{5,7,9,6}\}$

Out[*]=

pdf

True

tex


\end{frame}

Reidemeister 1

tex

\begin{frame}{\large Reidemeister 1}

[\ \def{p{=\tau_1^{1/2}} \def{m{=\tau_1^{-1/2}} \input{figs/R1.pdf_t} \]



pdf

In[*]:= $\{\mathcal{A}@\{X_{3,3,2,1}\} \equiv \tau_1^{-1/2} \mathcal{A}@\{P_{1,2}\}, \mathcal{A}@\{X_{1,2,3,3}\} \equiv \tau_1^{1/2} \mathcal{A}@\{P_{1,2}\},$
 $\mathcal{A}@\{\bar{X}_{1,3,3,2}\} \equiv \tau_1^{-1/2} \mathcal{A}@\{P_{1,2}\}, \mathcal{A}@\{\bar{X}_{3,1,2,3}\} \equiv \tau_1^{1/2} \mathcal{A}@\{P_{1,2}\}\}$

Out[*]=

pdf


{True, True, True, True}

tex

(So we have an invariant, up to rotation numbers).

\end{frame}

The Relation with the Multivariable Alexander Polynomial



tex

\begin{frame}{\large The Relation with the Multivariable Alexander Polynomial}

[\ \input{figs/Borromean.pdf_t} \]

pdf

$$\text{In[*]} := \text{MVA} = u^{-1/2} v^{-1/2} w^{-1/2} (u - 1) (v - 1) (w - 1);$$

pdf

$$\text{In[*]} := \mathbf{A} = \{ \bar{X}_{1,12,2,13} [u, v], \bar{X}_{13,2,6,3}, X_{8,4,9,3}, X_{4,10,5,9}, X_{6,17,7,16} [v, w], X_{15,8,16,7}, \bar{X}_{14,10,15,11}, \bar{X}_{11,17,12,14} \} // \mathcal{A} // \text{Last} // \text{Factor}$$

Out[*]=

pdf

$$\frac{(-1 + u)^2 (-1 + v) (-1 + w) (\text{Wedge}[] - x_5 \wedge \xi_1)}{u v}$$

pdf

$$\text{In[*]} := \mathbf{A} == u^{-1/2} (u - 1) u^0 v^{-1/2} w^{1/2} \text{MVA} (\text{Wedge}[] - x_5 \wedge \xi_1)$$

Out[*]=

pdf

True

tex

\end{frame}


Overcrossings Commute but Undercrossings don't

tex

\begin{frame}{\large Overcrossings Commute but Undercrossings don't}

tex

\[\input{figs/OUC.pdf_t} \]



pdf

$$\text{In[*]} := \mathcal{A} @ \{ X_{2,7,5,1}, X_{3,4,6,7} \} \equiv \mathcal{A} @ \{ X_{3,7,6,1}, X_{2,4,5,7} \}$$

Out[*]=

pdf

True

pdf

$$\text{In[*]} := \mathcal{A} @ \{ \bar{X}_{1,2,7,5}, \bar{X}_{7,3,4,6} \} \equiv \mathcal{A} @ \{ \bar{X}_{1,3,7,6}, \bar{X}_{7,2,4,5} \}$$

Out[*]=

pdf

False

tex

\end{frame}

The Conway Relation

tex

\begin{frame}{\large The Conway Relation\hfill (see~\cite{Conway:Enumeration})}

\[\input{figs/Conway.pdf_t} \]

pdf

$$In[*]:= \mathcal{A}@\{X_{2,3,4,1}[T, T]\} - \mathcal{A}@\{\bar{X}_{1,2,3,4}[T, T]\} \equiv (T^{-1/2} - T^{1/2}) \mathcal{A}@\{P_{1,4}[T], P_{2,3}[T]\}$$

Out[*]=
pdf

True

tex

```
\[ \includegraphics[height=36mm]{../../Projects/Gallery/Conway.png} \]
\end{frame}
```

Conway's Second Set of Identities

tex

```
\begin{frame}\large Conway's Second Set of Identities\hfill (see~\cite{Conway:Enumeration})
\[ \def\b{S=((u/v)^{1/2}+(u/v)^{-1/2})} \def{c}S=((u/v)^{1/2}+(u/v)^{-1/2})} \input{figs/Conway2nd.pdf_t}
\]
```

pdf

$$In[*]:= \mathcal{A}@\{X_{2,4,3,1}[v, u], X_{4,6,5,3}\} + \mathcal{A}@\{\bar{X}_{1,2,4,3}[u, v], \bar{X}_{3,4,6,5}\} \equiv (u^{1/2} v^{1/2} + u^{-1/2} v^{-1/2}) \mathcal{A}@\{P_{1,5}[u], P_{2,6}[v]\}$$

Out[*]=
pdf

True

pdf

$$In[*]:= \mathcal{A}@\{\bar{X}_{4,1,6,3}[v, u], \bar{X}_{3,2,5,4}\} + \mathcal{A}@\{X_{1,6,3,4}[u, v], X_{2,5,4,3}\} \equiv (u^{1/2} v^{-1/2} + u^{-1/2} v^{1/2}) \mathcal{A}@\{P_{1,5}[u], P_{2,6}[v]\}$$

Out[*]=
pdf


True

tex

```
\end{frame}
```

tex

```
\begin{frame}\null
{\bf Virtual versions} (Archibald,~\cite{Archibald:Thesis})
\[ \def{b}S=(\tau_1^{1/2}+\tau_1^{-1/2})} \def{c}S=(\tau_2^{1/2}+\tau_2^{-1/2})} \input{figs/Conway2ndV.pdf_t} \]
```



pdf

$$In[*]:= \mathcal{A}@\{X_{2,3,4,1}\} + \mathcal{A}@\{\bar{X}_{2,1,4,3}\} \equiv (\tau_1^{1/2} + \tau_1^{-1/2}) \mathcal{A}@\{P_{1,3}, P_{2,4}\}$$

Out[*]=
pdf

True

pdf

$$In[*]:= \mathcal{A}@\{\bar{X}_{1,2,3,4}\} + \mathcal{A}@\{X_{1,4,3,2}\} \equiv (\tau_2^{1/2} + \tau_2^{-1/2}) \mathcal{A}@\{P_{1,3}, P_{2,4}\}$$

Out[*]=
pdf

True


tex

\end{frame}

Conway's Third Identity

tex

\begin{frame}{\large Conway's Third Identity \hfill (see~\cite{Conway:Enumeration})}
 \[\input{figs/C3.pdf_t} \]



pdf

$$In[*]:= \mathcal{A}@\{X_{6,4,9,1}, \bar{X}_{4,5,7,8}, \bar{X}_{2,3,5,6}\} + \mathcal{A}@\{X_{2,4,5,1}, \bar{X}_{4,3,7,6}, X_{6,8,9,5}\} \equiv \mathcal{A}@\{\bar{X}_{1,6,4,9}, X_{5,7,8,4}, X_{3,5,6,2}\} + \mathcal{A}@\{\bar{X}_{1,2,4,5}, X_{3,7,6,4}, \bar{X}_{5,6,8,9}\}$$

Out[*]=
pdf


True

tex

\end{frame}

tex

\begin{frame}\null
 {\bf Virtual version} (Archibald,~\cite{Archibald:Thesis})
 \[\input{figs/C3V.pdf_t} \]



pdf

$$In[*]:= \mathcal{A}@\{X_{3,7,6,1}, \bar{X}_{7,2,4,5}\} + \mathcal{A}@\{X_{2,4,7,1}, X_{3,5,6,7}\} \equiv \mathcal{A}@\{X_{3,7,6,2}, X_{7,4,5,1}\} + \mathcal{A}@\{\bar{X}_{1,2,7,5}, X_{3,4,6,7}\}$$

Out[*]=


pdf

True

tex

\end{frame}

Jun Murakami's Fifth Axiom



tex

\begin{frame}{\large Jun Murakami's Fifth Axiom \hfill (see~\cite{MurakamiJ:StateModel})}
 \[\def\prop{\\$=\frac{\sqrt{S}(1-T)}{\sqrt{T}}\\$} \input{figs/MJ5.pdf_t} \]


pdf

$$In[*]:= \mathcal{A}@\{X_{1,4,2,5}[T, S], X_{4,3,5,2}\} \equiv \frac{\sqrt{S} (1 - T)}{\sqrt{T}} \mathcal{A}@\{P_{1,3}[T]\}$$

Out[*]=

pdf

True



tex

\[\includegraphics[height=20mm]{../Projects/Gallery/MurakamiJ.jpg} \]
 \end{frame}

tex

\begin{frame}\null
 {\bf Virtual versions} (Archibald,~\cite{Archibald:Thesis})
 \[\def\prop{\\$(T^{-1/2}-T^{1/2})\\$} \input{figs/MJ5V.pdf_t} \]

```
pdf
In[ ]:= A@{X3,2,3,1[S, T]} ≡ (T-1/2 - T1/2) A@{P1,2[T]}
```


```
Out[ ]=
pdf
True
```

```
pdf
In[ ]:= A@{X1,3,2,3}
Out[ ]=
pdf
A[{1}, {2}, <|ξ1 → τ1, x2 → τ1|>, 0]
```

```
tex
\end{frame}
```

Jun Murakami's Third Axiom

```
tex
\begin{frame}{\large Jun Murakami's Third Axiom\hfill(see~\cite{MurakamiJ:StateModel})}
\[\scalebox{0.64}{\input{figs/Murakami3.pdf_t}}\]
```



```
pdf
In[ ]:= A2112 = A@{X3,8,7,2, X7,10,9,1, X10,11,4,9, X8,6,5,11};
A1221 = A@{X2,8,7,1, X3,10,9,8, X10,6,11,9, X11,5,4,7};
A2211 = A@{X3,8,7,2, X8,6,9,7, X9,11,10,1, X11,5,4,10};
A1122 = A@{X2,8,7,1, X8,9,4,7, X3,11,10,9, X11,6,5,10};
A11 = A@{X2,8,7,1, X8,5,4,7, P3,6}; A22 = A@{X3,8,7,2, X8,6,5,7, P1,4};
A0 = A@{P1,4, P2,5, P3,6};
g+[z_] := z1/2 + z-1/2; g-[z_] := z1/2 - z-1/2;
g+[τ1] g-[τ2] A2112 - g-[τ2] g+[τ3] A1221 - g-[τ3 / τ1] (A2211 + A1122) +
g-[τ2 τ3 / τ1] g+[τ3] A11 - g+[τ1] g-[τ1 τ2 / τ3] A22 ≡ g-[τ32 / τ12] A0
```


```
Out[ ]=
pdf
True
```

```
tex
\end{frame}
```

The Naik-Stanford Double Delta Move

```
tex
\begin{frame}{\large The Naik-Stanford Double Delta Move\hfill(see~\cite{NaikStanford:Move})}
\vskip -5mm
```

```
\[
\includegraphics[height=20mm]{../../Projects/Gallery/Naik.jpg}
\qquad\scalebox{0.8}{\input{figs/NaikStanford.pdf_t}}\qquad
\includegraphics[height=20mm]{../../Projects/Gallery/Stanford.jpg}
\]
```



pdf

```
In[*]:= Timing[ $\mathcal{A} @ \{X_{6,10,28,24}[w, v], \bar{X}_{28,3,29,19}[w, v], X_{26,20,27,19}[w, v], \bar{X}_{27,23,11,24}[w, v],$   

 $X_{1,12,13,30}[u, w], \bar{X}_{13,5,14,25}[u, w], X_{17,26,18,25}[u, w], \bar{X}_{18,29,8,30}[u, w],$   

 $X_{4,7,22,15}[v, u], \bar{X}_{22,2,23,16}[v, u], X_{20,17,21,16}[v, u], \bar{X}_{21,14,9,15}[v, u]\} \equiv$   

 $\mathcal{A} @ \{X_{5,9,25,21}[w, v], \bar{X}_{25,4,26,22}[w, v], X_{29,23,30,22}[w, v], \bar{X}_{30,20,12,21}[w, v],$   

 $X_{2,11,16,27}[u, w], \bar{X}_{16,6,17,28}[u, w], X_{14,29,15,28}[u, w], \bar{X}_{15,26,7,27}[u, w],$   

 $X_{3,8,19,18}[v, u], \bar{X}_{19,1,20,13}[v, u], X_{23,14,24,13}[v, u], \bar{X}_{24,17,10,18}[v, u]\}$ 
```

Out[*]=
pdf

{190.422, True}

tex


```
\end{frame}
```

tex

```
\begin{frame}\null
{\bf Virtual Version 1} (Archibald,~\cite{Archibald:Thesis})
```

tex

```
\[ \input{figs/VNS1.pdf_t} \]
```



pdf

$$\text{In}[*]:= \mathcal{A}@\{X_{1,8,11,3}[u, v], \bar{X}_{11,2,12,7}[u, v], X_{12,10,13,4}[u, w], \bar{X}_{13,5,6,9}[u, w]\} \equiv \\ \mathcal{A}@\{X_{1,10,11,4}[u, w], \bar{X}_{11,5,12,9}[u, w], X_{12,8,13,3}[u, v], \bar{X}_{13,2,6,7}[u, v]\}$$

Out[*]=

pdf

True

tex


\end{frame}

tex

\begin{frame}\null
{\bf Virtual Version 2} (Archibald,~\cite{Archibald:Thesis})

tex

\[\input{figs/VNS2.pdf_t} \]



pdf

$$\text{In}[*]:= \mathcal{A}@\{\bar{X}_{20,1,10,13}[v, u], X_{3,14,19,13}[v, u], X_{14,11,15,21}[u, w], \bar{X}_{15,6,7,22}[u, w], \\ X_{2,12,16,22}[u, w], \bar{X}_{16,5,17,21}[u, w], \bar{X}_{19,17,9,18}[v, u], X_{4,8,20,18}[v, u]\} \equiv \\ \mathcal{A}@\{X_{1,11,13,21}[u, w], \bar{X}_{13,6,14,22}[u, w], \bar{X}_{20,14,10,15}[v, u], X_{3,7,19,15}[v, u], \\ \bar{X}_{19,2,9,16}[v, u], X_{4,17,20,16}[v, u], X_{17,12,18,22}[u, w], \bar{X}_{18,5,8,21}[u, w]\}$$

Out[*]=

pdf

True

tex

\end{frame}