http://drorbn.net/24-327 Dror Bar-Natan: Classes: 2024-25:

term-test-938a9

#1 Page 1 of 22



Do not open this notebook until instructed.

MAT 327 Introduction to Topology

Term Test

University of Toronto, October 16, 2024

Solve all 5 problems on this booklet.

The problems are of equal weight. You have an hour and fifty minutes to write this test.

	<u> </u>			<u> </u>	ļ	ļ	ļ	<u> </u>	<u> </u>	<u> </u>	ļ	<u> </u>	<u> </u>	<u> </u>
Lact	name													
цавс	Trance			_				1	_					
Stud	ent I	D num	mber											
	Ì							Ì				Ì	Ì	
	1	l .									l			

Notes

- No outside material allowed other than stationery, minimal hydration and snacks, and stuffed animals.
- Write your solution of each problem on the problem page and on the back of the problem page. If you run out of space you may continue into the scratch pages, but you **must** indicate this on the problem page or else the scratch pages will not be read.
- Neatness counts! Language counts! The *ideal* written solution to a problem looks like a page from a textbook; neat and clean and consisting of complete and grammatical sentences. Definitely phrases like "there exists" or "for every" cannot be skipped. Lectures are mostly made of spoken words, and so the blackboard part of proofs given during lectures often omits or shortens key phrases. The ideal written solution to a problem does not do that.

Good Luck!



term-test-938a9

#1 Page 2 of 22

term-test-938a9

22E5D538-5933-45AF-86B6-8A46323F3D1E

#1 Page 3 of 22


Neatness counts! Language counts!

Problem 1. Let *X* be a topological space and let $f, g: X \to \mathbb{R}$ be a pair of continuous functions.

- 1. Show that $\{x \in X : f(x) = g(x)\}\$ is a closed set in X.
- 2. A subset $D \subset X$ is called *dense* if it has a non-empty intersection with every closed set in X (example: $D = \mathbb{Q}$ is dense in $X = \mathbb{R}$). With f and g as before, show that if f is equal to g on a dense set D, then f is equal to g everywhere.

Tip. Don't start working! Read the whole exam first. You may wish to start with the questions that are easiest for you.

Your solution of Problem 1.





term-test-938a9 #1 Page 4 of 22

Neatness counts! Language counts!

Problem 2. Show that if a topological space X is Hausdorff and if x_1, \ldots, x_n are distinct points in X, then there exist open sets U_1, \ldots, U_n in X such that $\forall i \ x_i \in U_i$ and such that $\forall i \ \neq j, \ U_i \cap U_j = \emptyset$.

Tip. In math exams, "show" means "prove".


Your solution of Problem 2.

A9EEE175-D37E-4966-BEDA-5723EF422376

term-test-938a9

#1 Page 5 of 22







term-test-938a9
#1 Page 6 of 22

Neatness counts! Language counts!

Problem 3. Let X be a metrizable space and let $A \subset X$ be some subset of X. Show that a point x belongs to \bar{A} if and only if there exists a sequence a_i of points in A such that $a_i \to x$.

6FA2C00A-6CE7-4AB2-AE62-1F59BC886BE9


term-test-938a9

#1 Page 7 of 22



Tip. "If and only if" always means that there are two things to prove.

Your solution of Problem 3.





term-test-938a9 #1 Page 8 of 22

Neatness counts! Language counts!

Problem 4. Let X_n be a sequence of non-empty topological spaces whose topology is not the trivial topology. Show that the boxes topology on $X := \prod_{n=1}^{\infty} X_n$ is strictly stronger than the cylinders topology on X.

1AF45140-7F5F-47DD-AA36-EF3A911B9CF2

term-test-938a9

#1 Page 9 of 22



Tip. Here and almost always, a concise yet precise solution is better than a lengthy roundabout one.

Your solution of Problem 4.





term-test-938a9 #1 Page 10 of 22

Neatness counts! Language counts!

Problem 5. Let X be a topological space, let \sim be an equivalence relation on X, let $Y = X/\sim$ be the quotient set, and let $\pi: X \to Y$ be the natural projection.

61EE73E7-5D43-4D15-BF9C-30391B279BBF

term-test-938a9

#1 Page 11 of 22




- 1. Write the "functional" definition of the quotient topology on *Y*. Namely, list some functions or some families of functions whose range is *Y* or whose domain is *Y* that must be continuous relative to the quotient topology on *Y*.
- 2. Prove that if a quotient topology on *Y* exists, then it is unique. (Note that you are not asked to prove that the quotient topology exists!).

Tip. For all problems, you may want to start by writing "draft solutions" on the last pages of this notebook and only then write the perfected versions in the space allocated for the solutions.

Tip. Once you have finished writing an exam, if you have time left, it is always a good idea to go back and re-read and improve everything you have written, and perhaps even completely rewrite any parts that came out messy.

Your solution of Problem 5.





term-test-938a9 #1 Page 12 of 22

FD5663C0-D939-431E-A9C5-27445AA03EC0

term-test-938a9

#1 Page 13 of 22





term-test-938a9 #1 Page 14 of 22

81D6845F-4641-485A-AAB8-B65F1AB34593

term-test-938a9

#1 Page 15 of 22





term-test-938a9 #1 Page 16 of 22

C4890B3B-800A-412D-92D5-397A04C02AB3

term-test-938a9

#1 Page 17 of 22





term-test-938a9

#1 Page 18 of 22

A59E2BDA-DA0D-4385-B9B4-DF9958E2BC48

term-test-938a9

#1 Page 19 of 22





term-test-938a9 #1 Page 20 of 22

88F82E52-8130-4F1F-888E-2499A71C918A

term-test-938a9

#1 Page 21 of 22



