

Table[

CF[R[1, 2]i /. {gg_{2|4|5|7}[_] → 0, gg₆[x_] := $\frac{2-x}{2x^2}$,
gg₈[x_] := 1/x, cc₁ → 0}], {i, 2}] // AForm

$$\left\{ \begin{aligned} & \text{UU}\left[a[1, 1, \infty] + \text{aa}\left[-\frac{-1+e^{-b_1+b_1}}{b_1^2}, 1, 2, 1, \infty\right] + \right. \\ & \left. \text{aa}\left[\frac{1}{b_2}, 1, 2, 2, \infty\right] + \text{ca}\left[\frac{1-e^{-b_1}}{b_1}, \infty, 1, 2\right]\right], \\ & \text{UU}\left[a[e^{b_1}, 2, \infty] + a\left[-\frac{(-1+e^{b_1})b_2}{b_1}, 1, \infty\right] + \right. \\ & \left. \text{aa}\left[\frac{e^{-b_1}(-1+e^{b_1})^2}{b_1^2}, 1, 2, 1, \infty\right] + \text{aa}\left[\frac{1-e^{b_1}}{b_1 b_2}, 1, 2, 2, \infty\right] + \right. \\ & \left. \text{ao}\left[\frac{-1+e^{b_1}-e^{b_1}b_2}{b_1}, 1, \infty\right] + \text{ca}\left[-\frac{1-e^{-b_1}}{b_1}, \infty, 1, 2\right]\right] \end{aligned} \right\}$$